arXiv:1108.2310v1 [math.AG] 11 Aug 2011

ON G/N-HILB OF N-HILB

AKIRA ISHII, YUKARI ITO, AND ÁLVARO NOLLA DE CELIS

Abstract. In this paper we consider the iterated G-equivariant Hilbert scheme G/N-Hilb(N-Hilb) and prove that G/N-Hilb(N-Hilb($\mathbb{C}^3$)) is a crepant resolution of $\mathbb{C}^3/G$ isomorphic to the moduli space $\mathcal{M}_\theta(Q)$ of θ-stable representations of the McKay quiver Q for certain stability condition θ. We provide several explicit examples to illustrate this construction. We also consider the problem of when G/N-Hilb(N-Hilb) is isomorphic to G-Hilb showing the fact that these spaces are most of the times different.

Contents

1. Introduction 1
2. The moduli space of G-constellations and G/N-Hilb(N-Hilb) 3
3. The case G Abelian 7
4. Non-Abelian examples 11
5. When Hilb of Hilb is Hilb 19
References 25

1. Introduction

Let X be a nonsingular quasiprojective complex 3-fold and let $G \subset \operatorname{Aut} X$ be a finite subgroup such that the stabilizer subgroup of any point $x \in X$ acts on the tangent space T_xX as a subgroup of $\mathrm{SL}(T_xX)$. We have the following celebrated theorem of Bridgeland, King and Reid:

Theorem 1.1 ([BKR]). *$Y = G$-Hilb(X) is irreducible and $f : Y \to X/G$ is a crepant resolution. Furthermore, $\Phi : \mathrm{D}(G\text{-Hilb}(X)) \longrightarrow \mathrm{D}^G(X)$ is an equivalence of derived categories.*

Our framework is the following: let $G \subset \mathrm{SL}(3,\mathbb{C})$ finite and $N \trianglelefteq G$ a normal subgroup. First consider the action of N on $\mathbb{C}^3$ and take the crepant resolution $Y := N$-Hilb($\mathbb{C}^3$). Finally, act with G/N on Y to obtain G/N-Hilb(Y).

$$\begin{tikzcd}
G/N\text{-Hilb}(Y) \arrow[dr, "\tau_2"'] & & Y = N\text{-Hilb}(\mathbb{C}^3) \arrow[dl, "\pi_2"] \arrow[dr, "\tau_1"'] & & \mathbb{C}^3 \arrow[dl, "\pi_1"] \\
& Y/(G/N) & & \mathbb{C}^3/N &
\end{tikzcd}$$

As an immediate consequence of Theorem 1.1 we have the following corollary.

Corollary 1.2. *G/N-Hilb(N-Hilb($\mathbb{C}^3$)) is a crepant resolution of $\mathbb{C}^3/G$.*

Date: August 12, 2011.
The first author are supported by JSPS grant No.21540039 and No. 23540045. The second authors are supported by JSPS grant No.23540045, No.22224001 and No.09F09768. The third author is supported by FY2009 JSPS Fellowship for Foreign Researchers and JSPS grant No.09F09768.

In general G-Hilb($\mathbb{C}^3$) and G/N-Hilb(N-Hilb($\mathbb{C}^3$)) are non-isomorphic quasiprojective varieties, and even when they coincide, as moduli spaces of representations of the McKay quiver almost always they belong to different chambers in the space of stability conditions Θ. Nevertheless, since any two crepant resolutions of a three-dimensional projective variety Y with only terminal singularities are derived equivalent by [Bri], we have that $\mathrm{D}^b(G\text{-Hilb}(\mathbb{C}^3)) \cong \mathrm{D}^b(G/N\text{-Hilb}(N\text{-Hilb}(\mathbb{C}^3)))$.

The similar construction was considered before by the second author in [Ito94], [Ito95] in the case of trihedrals subgroups in $\mathrm{SL}(3,\mathbb{C})$. The trihedral group is a non-Abelian finite subgroup generated by diagonal matrices and a matrix $T := \left(\begin{smallmatrix} 0&1&0 \\ 0&0&1 \\ 1&0&0 \end{smallmatrix}\right)$. Let us change the name of the groups, $G/T = N$. Then in Ito's construction, we require that G/N acts on the crepant resolution Y of $\mathbb{C}^3/N$ symmetrically on the exceptional locus. Therefore, this construction gives the G/N-Hilb(N-Hilb($\mathbb{C}^3$)) when $Y = N$-Hilb($\mathbb{C}^3$).

This construction can be extended in a natural way to obtain crepant resolutions of $\mathbb{C}^3/G$ for any finite non-simple group $G \subset \mathrm{SL}(3,\mathbb{C})$ (see [YY] for a classification of such groups). In general, if we consider the filtration of groups $\{1\} = N_k \trianglelefteq N_{k-1} \trianglelefteq \ldots \trianglelefteq N_1 \trianglelefteq N_0 = G$ where every subgroup N_i is normal in N_{i+1}, then the iterated Hilbert scheme described here is crepant. In particular, it is always possible to find such a crepant resolution with N_i/N_{i+1} Abelian subgroups for every i.

Denote by $\mathrm{Irr}(G)$ the set of irreducible representations of G and let (Q,R) be the McKay quiver of G with relations R. For $\mathbf{d} = (\dim(\rho))_{\rho\in\mathrm{Irr}(G)}$ and any generic θ in the space of stability conditions Θ we can define $\mathcal{M}_{\theta,\mathbf{d}}(Q,R)$ to be the moduli space of θ-stable representations of Q. Moreover, there exists a chamber $C \in \Theta$ in which the GIT quotient $\mathcal{M}_{\theta,\mathbf{d}}(Q,R)$ is constant for all $\theta \in C$ (see [Tha]). Thus we may also denote this moduli space simply by $\mathcal{M}_C$. The methods in [BKR] can be applied to prove that $\tau : \mathcal{M}_C \longrightarrow \mathbb{C}^3/G$ is a crepant resolution and $\Phi_C : D(\mathcal{M}_C) \longrightarrow \mathrm{D}^G(\mathbb{C}^3)$ is an equivalence of categories (see [CI] Section 2).

For these moduli spaces, the problem of whether every (projective) crepant resolution of $\mathbb{C}^3/G$ is a moduli of representations of the McKay quiver was treated by Craw and the first author in the case of Abelian group actions.

Theorem 1.3 ([CI]). *For a finite Abelian subgroup $A \subset \mathrm{SL}(3,\mathbb{C})$ let $Y \to \mathbb{C}^3/A$ be a projective crepant resolution. Then $Y \cong \mathcal{M}_C$ for some chamber $C \subset \Theta$.*

Then Craw and Ishii proposed the following conjecture.

Conjeture 1.4. *For a finite subgroup $G \subset \mathrm{SL}(3,\mathbb{C})$ let $Y \to \mathbb{C}^3/G$ be a projective crepant resolution. Then $Y \cong \mathcal{M}_C$ for some chamber $C \subset \Theta$.*

In this paper, we consider the G/N-Hilb(N-Hilb($\mathbb{C}^3$)) and show that it is not only a crepant resolution, but also a moduli space for a particular chamber $C \subset \Theta$ as follows:

Theorem 1.5. *(= 2.6) Let $G \subset \mathrm{SL}(3,\mathbb{C})$ finite and let $N \trianglelefteq G$ be a normal subgroup. The crepant resolution G/N-Hilb(N-Hilb($\mathbb{C}^3$)) is isomorphic to a moduli space of G-constellations $\mathcal{M}_C$ for some chamber $C \subset \Theta$.*

Thus our main result shows that the conjecture holds for the family G/N-Hilb(N-Hilb($\mathbb{C}^3$)) of crepant resolutions for general $G \subset \mathrm{SL}(3,\mathbb{C})$. Then it is natural to ask how much different G/N-Hilb(N-Hilb($\mathbb{C}^3$)) is from G-Hilb($\mathbb{C}^3$). In this paper, we consider consider the problem of when the iterated Hilbert scheme G/N-Hilb(N-Hilb($\mathbb{C}^3$)) is isomorphic to G-Hilb($\mathbb{C}^3$). We give a complete answer for this problem as moduli spaces in general, and we also show when they are isomorphic as algebraic varieties when the group G is Abelian. Indeed, these varieties are different in most cases. Such a statement in non-Abelian cases would be an (small) evidence for Conjecture 1.4.

This paper is organized as follows. In section 2, we introduce the moduli space of G-constellations and see the stability for G/N-Hilb(N-Hilb($\mathbb{C}^3$)) to prove Theorem1.5. In section 3, we show some examples of G/N-Hilb(N-Hilb($\mathbb{C}^3$)) when G is Abelian. The construction of crepant resolutions for Abelian quotient singularities are well known as toric resolutions where one of them is the G-Hilb($\mathbb{C}^3$). However, if we have several choices for normal subgroups N of G, then we have several G/N-Hilb(N-Hilb($\mathbb{C}^3$)) which can be obtained by a finite sequence of flops from G-Hilb($\mathbb{C}^3$). Moreover, the actions of G/N on N-Hilb($\mathbb{C}^3$) are also interesting. In section 4, we give several examples of G/N-Hilb(N-Hilb($\mathbb{C}^3$)) when G is non-Abelian. We also describe the G-constellations and McKay quiver with relations. In section 5, we show when G/N-Hilb(N-Hilb($\mathbb{C}^3$)) is isomorphic to G-Hilb($\mathbb{C}^3$) as moduli spaces and as algebraic varieties.

The authors thank Alastair King and Laurent Demonet for their valuable comments and Alastair Craw and Michael Wemyss for useful discussions.

2. THE MODULI SPACE OF G-CONSTELLATIONS AND G/N-HILB(N-HILB)

Recall that G-Hilb($\mathbb{C}^n$) is the moduli space of G-clusters, where a *G-cluster* $\mathcal{Z}$ is a G-invariant subscheme $\mathcal{Z} \subset \mathbb{C}^n$ such that $H^0(\mathcal{O}_Z) \cong \mathbb{C}[G]$ the regular representation of G as $\mathbb{C}[G]$-modules. Thus a point $y \in G/N$-Hilb(N-Hilb($\mathbb{C}^3$)) is a G/N-cluster of N-clusters.

The first observation that appears is that y may not be a G-cluster. Therefore, in order to construct the moduli space of such objects we need the generalized notion of G-cluster called *G-constellation* (see [CI] and [Cra01], Chapter 5).

Definition 2.1. A *G-constellation* $\mathcal{F}$ on X is a G-equivariant coherent sheaf on X such that $H^0(\mathcal{F}) \cong R_G$ as $\mathbb{C}[G]$-modules.

Notice that a G-cluster is a G-equivariant the $\mathbb{C}[x_1,\dots,x_n]$-module $\mathcal{O}_{\mathcal{Z}}$ generated from 1 mod $I_{\mathcal{Z}}$, which is precisely a G-constellation generated form ρ_0.

Equivalently, when $X = \mathbb{C}^n$ a G-constellation $\mathcal{F}$ on X is a representation of the McKay quiver Q (see [CI]). We consider both interpretations of G-constellations indistinctly throughout the paper.

We fix the following notation in this paper. Denote by $Y_1 := N$-Hilb($\mathbb{C}^n$) and $Y_2 := G/N$-Hilb(Y_1). Then we have the diagram

$$\begin{tikzcd}
& \mathcal{Z}_2 \arrow[dl, "p_2"'] \arrow[dr, "q_2"] & & \mathcal{Z}_1 \arrow[dl, "p_1"'] \arrow[dr, "q_1"] & \\
Y_2 \arrow[dr, "\tau_2"'] & & Y_1 \arrow[dl, "\pi_2"] \arrow[dr, "\tau_1"'] & & \mathbb{C}^n \arrow[dl, "\pi_1"] \\
& Y_1/(G/N) & & \mathbb{C}^n/N &
\end{tikzcd}$$

where $\mathcal{Z}_1$ and $\mathcal{Z}_2$ are the universal families for Y_1 and Y_2 respectively.

Lemma 2.2. *Every point in the connected component of G/N-Hilb(N-Hilb($\mathbb{C}^n$)) dominating $\mathbb{C}^3/G$ is a G-constellation on $\mathbb{C}^n$. More precisely, there is a canonically defined flat family of G-constellations parametrised by this connected component G/N-Hilb(N-Hilb($\mathbb{C}^n$)).*

Proof. Consider the fibre product $\mathcal{Z}_2 \times_{Y_1} \mathcal{Z}_1 \subset Y_2 \times Y_1 \times \mathbb{C}^n$ and the projection $p_{02} : Y_2 \times Y_1 \times \mathbb{C}^n \longrightarrow Y_2 \times \mathbb{C}^n$ onto the first and third factor. Then $p_{02*}(\mathcal{O}_{\mathcal{Z}_2 \times_{Y_1} \mathcal{Z}_1})$ is a G-equivariant coherent sheaf on $Y_2 \times \mathbb{C}^n$, flat over Y_2. For a closed point $y \in Y_2$, let $\mathcal{O}_W = q_{2*}p_2^*\mathcal{O}_y \subset Y_1$ be the corresponding G/N-cluster. Then the fibre of $p_{02*}(\mathcal{O}_{\mathcal{Z}_2 \times_{Y_1} \mathcal{Z}_1})$ over y is $q_{1*}p_1^*\mathcal{O}_W$. Especially, if y lies over a free orbit in $\mathbb{C}^n/G$, then $q_{1*}p_1^*\mathcal{O}_W$ is the G-cluster supported by the free orbit and it is the regular representation as a G-module. Since $p_{02*}(\mathcal{O}_{\mathcal{Z}_2 \times_{Y_1} \mathcal{Z}_1})$ is flat over Y_2, it is a flat family of G-constellations in the connected component containing free orbits.

□

2.1. **Stability for G/N-Hilb(N-Hilb($\mathbb{C}^3$)).** Let $\Theta = \Theta^G := \{\theta \in \mathrm{Hom}_{\mathbb{Z}}(R[G], \mathbb{Q}) | \theta(R_G) = 0\}$ with $R[G]$ the representation ring. The notion of stability by [Kin] translates into the language of G-constellations as follows. For $\theta \in \Theta$, a G-constellation $\mathcal{F}$ is θ-(semi)stable if $\theta(\mathcal{E}) = \theta(H^0(\mathcal{E})) > 0 = \theta(\mathcal{F})$ for $0 \subsetneq \mathcal{E} \subsetneq \mathcal{F}$ (with the usual $\geq$ for semistability). With quiver-theoretic point of view, if M is a representation of Q of dimension vector $\mathbf{d} = (d_i)_{i \in Q_0}$ then the notion of stability for M is given as follows: let $\theta \in \mathbb{Q}^{Q_0}$ and define $\theta(M) := \sum \theta_i d_i$. Then M is θ-(semi)stable if $\theta(M') > 0 = \theta(M)$ for $0 \subsetneq M' \subsetneq M$ (with the usual $\geq$ for semistability).

It is known from [IN] that G-Hilb($\mathbb{C}^3$) can be considered as a moduli $\mathcal{M}_\theta$ of θ-stable representations of the McKay quiver of G for any stability condition $\theta \in \Theta$ satisfying $\theta(\rho) > 0$ for every non-trivial irreducible representation ρ of G (and hence $\theta(\rho_0) < 0$ for the trivial representation ρ_0), also called ***0**-generated* stability.

Remark 2.3. In general, the chamber of stability parameters defining G-Hilb is larger than the cone defined by the inequalities above. In the following argument, θ^N and $\theta^{G/N}$ can be chosen arbitrarily in the chambers of N-Hilb and G/N-Hilb respectively.

Let $\theta^N \in \Theta^N$ and $\theta^{G/N} \in \Theta^{G/N}$ be ***0**-generated* stabilities for N and G/N respectively.

Definition 2.4. Let $\rho \in \mathrm{Irr}(G)$ and $\theta \in \Theta$. We define

$$\theta(\rho) := \begin{cases} \theta^N(\rho|_N) + \varepsilon \cdot \theta^{G/N}(\rho) & \text{if } \rho \in \mathrm{Irr}(G/N) \\ \theta^N(\rho|_N) & \text{if } \rho \notin \mathrm{Irr}(G/N) \end{cases}$$

where $0 < \varepsilon << 1$.

Note that $\theta^N \in \Theta^N$ can be regarded as an element of Θ via the restriction map $R(G) \to R(N)$. The condition $\rho \in \mathrm{Irr}(G/N)$ as a representation of G means that ρ is trivial for every element in N. It is straightforward to check that $\theta(R_G) = 0$ for the regular representation R_G as required.

From now on we restrict to the case $G \subset \mathrm{SL}(3, \mathbb{C})$. Consider the functor

$$\Phi : D^b(\mathrm{Coh}^{G/N} Y_1) \to D^b(\mathrm{Coh}^G \mathbb{C}^3)$$

defined by $\Phi(-) = \mathbb{R}q_{1*}p_1^*(-)$. Then it is an equivalence of triangulated categories by [BKR](see also [IU, Theorem 3.1]). Let $\mathrm{Coh}_0^{G/N}(Y_1)$ denote the Abelian category of G/N-equivariant coherent sheaves on Y_1 with zero-dimensional supports. Then Φ sends objects of $\mathrm{Coh}_0^G(Y_1)$ to G-equivariant sheaves with zero-dimensional supports and Φ is exact on $\mathrm{Coh}_0^G(Y_1)$.

Lemma 2.5. *Let $G \subset \mathrm{SL}(3, \mathbb{C})$ be a finite subgroup and N be a normal subgroup of G. Then for an object $E \in \mathrm{Coh}_0^{G/N} Y_1$, $\Phi(E) \in \mathrm{Coh}^G \mathbb{C}^3$ is θ^N-semistable. Moreover, if $\mathcal{F} \subseteq \Phi(E)$ is a G-equivariant subsheaf of $\Phi(E)$ with $\theta^N(\mathcal{F}) = 0$, then there is a G/N-equivariant subsheaf F of E such that $\mathcal{F} = \Phi(F)$.*

Proof. Let $F_0 : D^b(\mathrm{Coh}^G \mathbb{C}^3) \to D^b(\mathrm{Coh}^N \mathbb{C}^3)$ and $F_1 : D^b(\mathrm{Coh}^{G/N} Y_1) \to D^b(\mathrm{Coh}\, Y_1)$ be the forgetful functors. We have a commutative diagram

$$\begin{tikzcd} D^b(\mathrm{Coh}^{G/N} Y_1) \arrow[r, "\Phi", "\sim"'] \arrow[d, "F_1"'] & D^b(\mathrm{Coh}^G \mathbb{C}^3) \arrow[d, "F_0"] \\ D^b(\mathrm{Coh}\, Y_1) \arrow[r, "\sim", "\Phi^N"'] & D^b(\mathrm{Coh}^N \mathbb{C}^3) \end{tikzcd}$$

where Φ^N is the functor which is defined in the same way as Φ and is an equivalence by [BKR]. Since $F_1(E)$ has a filtration in $\mathrm{Coh}\, Y_1$ whose factors are skyscraper sheaves, $\Phi^N(F_1(E))$ has a filtration in $\mathrm{Coh}^N \mathbb{C}^3$ whose factors are N-clusters. Therefore, $F_0(\Phi(E)) \cong \Phi^N(F_1(E))$ is θ^N-semistable. Now for any G-invariant subsheaf $\mathcal{F}$ of $\Phi(E)$, we have $\theta^N(\mathcal{F}) = \theta^N(F_0(\mathcal{F})) \geq 0$ by the semistability of $F_0(\Phi(E))$, which shows that $\Phi(E)$ is θ^N-semistable.

Suppose $\mathcal{F} \subseteq \Phi(E)$ is a G-invariant subsheaf with $\theta^N(\mathcal{F}) = 0$. Since $F_0(\Phi(E))$ is a θ^N-semistable N-sheaf whose Jordan-Hölder factors are N-clusters, $F_0(\mathcal{F})$ is also θ^N-semistable and $F_0(\mathcal{F})$ also has a filtration in $\mathrm{Coh}^N(\mathbb{C}^3)$ whose factors are N-clusters, by the Jordan-Hölder theorem for semistable sheaves. Note that the equivalence Φ^N induces an isomorphism

$$\mathrm{Hom}(\mathcal{O}_y, F_1(E)) \cong \mathrm{Hom}(\Phi^N(\mathcal{O}_y), \Phi^N(F_1(E)))$$

for any closed point $y \in Y_1$. Then we can show that there is a unique subsheaf F' of $F_1(E)$ such that $F_0(\mathcal{F}) = \Phi^N(F')$, by the induction on the length of the filtration on $F_0(\mathcal{F})$. F' must be preserved by the action of G/N by its uniqueness, which shows that F' is of the form $F_1(F)$ for a G/N-invariant subsheaf $F \subseteq E$. $\square$

Theorem 2.6. *Let $\theta \in \Theta$ be as in Definition 2.4. Let $G \subset \mathrm{SL}(3, \mathbb{C})$ be a finite subgroup and N be a normal subgroup of G. Then*

$$G/N\text{-}Hilb(N\text{-}Hilb(\mathbb{C}^3)) \cong \mathcal{M}_C$$

for any generic stability condition in the chamber $C \subset \Theta$ which contains θ.

Proof. G/N-Hilb(N-Hilb($\mathbb{C}^3$)) parametrises a family of G-constellations of the form $\Phi(\mathcal{O}_W)$ where $W \subset Y_1$ is a G/N-cluster. Take a G-invariant subsheaf $\mathcal{F}$ of $\Phi(\mathcal{O}_W)$. If $\theta^N(\mathcal{F}) > 0$, then we have $\theta(\mathcal{F}) > 0$ by the assumption $\varepsilon << 1$ and we may assume $\theta^N(\mathcal{F}) = 0$. In this case, there is a G/N-invariant subsheaf $F \subseteq \mathcal{O}_W$ such that $\mathcal{F} = \Phi(F)$ by Lemma 2.5. Note that we have $\tau_{1*}F \cong (\pi_{1*}\mathcal{F})^N$ by the definition of Φ, which implies that the number of copies of an irreducible representation of G/N appearing in $H^0(F)$ is the same as that in $H^0(\mathcal{F})$, proving $\theta(\mathcal{F}) = \varepsilon\theta^{G/N}(F) > 0$. Thus we obtain the θ-stability of $\Phi(\mathcal{O}_W)$ and hence Φ induces a morphism $f : G/N\text{-Hilb}(N\text{-Hilb}(\mathbb{C}^3)) \to \mathcal{M}_{\theta,d}(Q)$. Now since $G \subset \mathrm{SL}(3, \mathbb{C})$ both are crepant resolutions of $\mathbb{C}^3/G$ so f is an isomorphism. $\square$

Remark 2.7. In the above proof, we can replace θ^N by an arbitrary G/N-invariant generic stability parameters for N-constellations. Especially, we have similar results for iterated constuctions such as "Hilb of Hilb of Hilb".

Note that to let $\theta^{G/N}$ be general, we have to define the moduli space of θ-stable G-constellations on a quasi-projective variety with G-action, which can be done by patching local constructions.

2.2. From $\mathrm{Irr}(N)$ to $\mathrm{Irr}(G)$.

We study how the irreducible representations of G can be obtained from irreducible representations of N.

Let $G \subset \mathrm{GL}(n, \mathbb{C})$ be a finite subgroup and let N be a normal subgroup of G. Let $\mathrm{Irr}(N) = \{\sigma_0, \ldots, \sigma_p\}$ be the set of irreducible representations of N with $d_i^N := \dim(\sigma_i)$ and denote by $I = \{0, \ldots, p-1\}$ the set of subindices. Let us also denote by $\mathrm{Irr}(G/N) = \{\tau_0, \ldots, \tau_{h_0}\}$ the set of irreducible representations of G/N with $d_j^{G/N} := \dim(\tau_j)$.

The group G/N acts on $\mathrm{Irr}\, N$ as follows: since N is normal, G acts on N by conjugation $g \cdot h := ghg^{-1}$, for $g \in G, h \in N$. Let $\sigma \in \mathrm{Irr}(N)$ and consider its character $\chi_\sigma : N \to \mathbb{C}$. Then G acts on the characters χ_σ by $g \cdot \chi_\sigma(h) := \chi_\sigma(g^{-1}hg)$. Since the character is a function on the conjugacy classes in N this action is constant in the cosets gN, i.e. $gh \cdot \chi_\sigma(x) = gh' \cdot \chi_\sigma(x) = \chi_\sigma(g^{-1}xg)$, for $g \in G$, $x, h, h' \in N$. Therefore G/N acts on the characters of N, which induces an action on $\mathrm{Irr}(N)$ by

$$g \cdot \rho = \sigma \iff g \cdot \chi_\rho = \chi_\sigma$$

for $\rho, \sigma \in \mathrm{Irr}(N)$, $g \in G/N$. In particular $g \cdot \chi_\rho(1) = \chi_\sigma(1) = \dim(\sigma)$, thus representations in the same orbit have the same dimension.

Choose a set of representatives of the classes in $\mathrm{Irr}\, N$ under the action of G/N and denote by $\tilde{I} = \{0, \ldots, k\} \subseteq I$ the corresponding subset of subindices. For any $i \in \tilde{I}$ consider the orbit $\mathrm{Orb}(\sigma_i)$ of σ_i under G/N of length n_i. Let G_i be the stabilizer and let $\mathrm{Irr}(G_i) = \{\tau_i^0, \ldots, \tau_i^{h_i}\}$ the set of irreducible representations of G_i. Recall that if σ_i and σ_j are in the same orbit then G_i

and G_j are conjugate, in particular isomorphic. The trivial representation $\sigma_0 \in \mathrm{Irr}(N)$ is always fixed so that $G_0 = G/N$ and $\tau_0^j = \tau_j$ for all j.

The irreducible representations of G are obtained as follows: for every $i \in \tilde{I}$ the representations in the orbit $\mathrm{Orb}(\sigma_i)$ combine to give $h_i + 1$ irreducible representations ρ_i^j for $j = 0, \dots, h_i$ with $\dim(\rho_i^j) = n_i d_i^N \dim(\tau_i^j)$ (see Table 1). In particular, if σ_i is fixed by G/N then it give rise to h_0+1 irreducible representations, where each of them corresponds to an irreducible representation of G/N. Note that ρ_0^0 is the trivial representation of G.

σ_0	$\mathrm{Orb}(\sigma_1)$	$\dots$	$\mathrm{Orb}(\sigma_k)$
ρ_0^0	ρ_1^0		ρ_k^0
ρ_0^1			
$\vdots$	$\vdots$	$\dots$	$\vdots$
	$\rho_1^{h_1}$		$\rho_k^{h_k}$
$\rho_0^{h_0}$			

TABLE 1. Irreducible representations of G from the action of G/N into $\mathrm{Irr}(N)$.

Recall that for any finite group G we have that $|G| = \sum_{\rho \in \mathrm{Irr}(G)} \dim(\rho)^2$ and if G acts on a set X then $|\operatorname{Orb}(x)| = |G|/|G_x|$ where G_x is the stabilizer subgroup of $x \in X$. Then

$$\sum_{i,j} (\dim \rho_i^j)^2 = \sum_{i=0}^{k} \sum_{j=0}^{h_i} (n_i d_i \dim \tau_i^j)^2 = |G/H| \sum_{i \in \tilde{I}} \frac{|G/H|}{|G_i|} d_i^2 = |G/H||H| = |G|.$$

Thus ρ_i^j are all the irreducible representations of G.

Remark 2.8. The action of G/N on $\mathrm{Irr}(N)$ to produce $\mathrm{Irr}(G)$ can be translated into the McKay quiver N, where every vertex corresponds to a irreducible representation of N. Then G/N acts on the set of vertices and on the set of arrows of Q, as well as on the path algebra $\Bbbk Q$ permuting the set of primitive idempotents $\{e_i | i \in I\}$. We thus can construct the McKay quiver of G as the G/N-orbifold quiver of the McKay quiver of N. See [Dem] for the general formulation and [NdC] for the case of binary dihedral groups in $\mathrm{GL}(2, \mathbb{C})$.

Let us describe the stability parameter defined in Definition 2.4 which is shown in Table 2. We are going to use the $\mathbf{0}$-generated stabilities for the groups N and G/N separately so let us denote them as follows:

$$\theta^N \in \mathbb{Q}^{p+1} \text{ such that } \theta_i^N := \theta^N(\sigma_i) > 0 \text{ for } i \neq 0 \text{ and } \sigma_i \in \mathrm{Irr}(N).$$

$$\theta^{G/N} \in \mathbb{Q}^{h_0+1} \text{ such that } \theta_j^{G/N} := \theta^{G/N}(\tau_j) > 0 \text{ for } j \neq 0 \text{ and } \tau_j \in \mathrm{Irr}(G/N).$$

In particular we have $\sum_{i=0}^{p} d_i^N \theta_i^N = 0$ and $\sum_{i=0}^{h_0} d_j^{G/N} \theta_j^{G/N} = 0$, so that

$$\theta_0^N = -\sum_{i=1}^{p} d_i^N \theta_i^N \text{ and } \theta_0^{G/N} = -\sum_{j=1}^{h_0} d_j^{G/N} \theta_j^{G/N}$$

σ_0	$\mathrm{Orb}(\sigma_1)$	$\dots$	$\mathrm{Orb}(\sigma_k)$
$\theta_0^N+\varepsilon\theta_0^{G/N}$	$\dim(\tau_1^0)\cdot\sum_{\sigma_i\in\mathrm{Orb}(\sigma_1)}\theta_i^N$	$\dots$	$\dim(\tau_k^0)\cdot\sum_{\sigma_i\in\mathrm{Orb}(\sigma_k)}\theta_i^N$
$d_1^{G/N}\theta_0^N+\varepsilon\theta_1^{G/N}$			
$\vdots$	$\vdots$		$\vdots$
$d_{h_0}^{G/N}\theta_0^N+\varepsilon\theta_{h_0}^{G/N}$	$\dim(\tau_1^{h_1})\cdot\sum_{\sigma_i\in\mathrm{Orb}(\sigma_1)}\theta_i^N$		$\dim(\tau_k^{h_k})\cdot\sum_{\sigma_i\in\mathrm{Orb}(\sigma_k)}\theta_i^N$

TABLE 2. Stability condition θ in terms of θ^N and $\theta^{G/N}$.

3. THE CASE G ABELIAN

Let $G\subset\mathrm{SL}(3,\mathbb{C})$ be an Abelian subgroup and let $A\lhd G$ be a normal subgroup of G with $|A|=p$ and $|G/A|=q$. After introducing the toric notations that are needed, we describe how to calculate the triangulation of the junior simplex Δ corresponding to $Y:=G/A\text{-Hilb}(A\text{-Hilb}(\mathbb{C}^3))$ and we construct explicitly every G-constellation in Y from the A-clusters. Then we describe a method to calculate the local coordinates of a moduli space of G-constellations using the McKay quiver and finish the section describing the stability condition in the Abelian case.

Every element of G can be written of the form $g=\mathrm{diag}(\varepsilon^{a_1},\varepsilon^{a_2},\varepsilon^{a_3})$ where ε is a rth primitive root of unity and $0\leq a_i<r$. Let $L\supset\mathbb{Z}^3$ the lattice generated by the elements of G written in the form $\frac{1}{r}(a_1,a_2,a_3)$ and let $M:=L^\vee$ the dual lattice of Laurent monomials. The *junior simplex* is the triangle $\Delta\subset L_\mathbb{R}:=L\otimes_\mathbb{Z}\mathbb{R}$ with vertices the standard basis e_1,e_2,e_3. We denote by $\mathbb{R}^2_\Delta$ the affine plane spanned by Δ and $\mathbb{Z}^2_\Delta:=L\cap\mathbb{R}^2_\Delta$. Recall that Δ contains all lattice points with $a_1+a_2+a_3=r$ and triangulations of Δ are in 1-to-1 correspondence with crepant resolutions of $\mathbb{C}^3/G$.

First consider the action of A on $\mathbb{C}^3$. In [CR] Craw and Reid give method to triangulate Δ into p regular triangles Δ_i which produces the crepant resolution $A\text{-Hilb}(\mathbb{C}^3)$. This triangulation shows that $A\text{-Hilb}(\mathbb{C}^3)\cong\bigcup_{i=1}^p Y_i$ where $Y_i:=\sigma(\Delta_i)\cong\mathbb{C}^3_{\alpha_i,\beta_i,\gamma_i}$ is the affine toric variety associated to the triangle Δ_i, and α_i, β_i, γ_i are Laurent monomials in x, y and z.

The action of G/A on $A\text{-Hilb}(\mathbb{C}^3)$ is again Abelian so it is given by diagonal matrices, thus it acts on every Y_i separately. For every triangle Δ_i with $i=1,\dots,p$, we form the the toric singular quotient $Y_i/(G/A)$ and take $G/A\text{-Hilb}(Y_i)$ as crepant resolution. Therefore,

$$G/A\text{-Hilb}(A\text{-Hilb}(\mathbb{C}^3))=\bigcup_{i=1}^p G/A\text{-Hilb}(Y_i)$$

In other words, the triangulation of Δ which gives $G/A\text{-Hilb}(A\text{-Hilb}(\mathbb{C}^3))$ is produced in two steps: Firstly calculate $A\text{-Hilb}(\mathbb{C}^3)$ according to [CR] to obtain $\Delta=\bigcup_{i=1}^p\Delta_i$. Secondly, triangulate every Δ_i into q regular triangles with the same method according to the $\mathbb{Z}/q$-action of G/A into Y_i to produce $\Delta=\bigcup\Delta_{ij}$ for $i=1,\dots,p$, $j=1,\dots,q$. Obviously, the same process of successive triangulations can be done as many times as nontrivial normal subgroups we have in a filtration of G. See Figure 1.

We now look at how $G/A\text{-Hilb}(A\text{-Hilb}(\mathbb{C}^3))$ can be constructed explicitly as a moduli of G-constellations. As an A-constellation, a point $\mathcal{F}\in A\text{-Hilb}(\mathbb{C}^3)$ is an A-equivariant coherent sheaf on $\mathbb{C}^3$ such that $H^0(\mathcal{F})\cong\mathbb{C}[A]\cong\bigoplus_{\sigma\in\mathrm{Irr}(A)}\sigma$, and $\mathcal{F}\cong\mathcal{O}_Z$ for some A-cluster Z. Therefore, locally at $U\subset A\text{-Hilb}(\mathbb{C}^3)$ we can take a basis $\Gamma:=\{\lambda_\sigma|\sigma\in\mathrm{Irr}(A)\}$ of $H^0(\mathcal{F})$ to be an A-graph.


$\mathbb{Z}/2$-Hilb $\mathbb{Z}/3$-Hilb($\mathbb{Z}/2$-Hilb) $\mathbb{Z}/5$-Hilb($\mathbb{Z}/3$-Hilb($Z/2$-Hilb))

FIGURE 1. Successive triangulations of Δ for a group G of order $r = 30 = 2{\cdot}3{\cdot}5$.

That is, λ_σ is a monomial in $\mathbb{C}[x,y,z]$ and if $x^iy^jz^k \in \Gamma$ then $x^{i'}y^{j'}z^{k'} \in \Gamma$ for any $i' \leq i$, $j' \leq j$ and $k' \leq k$ (see [Nak]). We call Γ the *building block* for the open set U.

It is also known from [Nak] that $U = \sigma(\Delta_i) \cong \mathbb{C}^3_{a,b,c}$ where $a = \frac{f_\sigma}{\lambda_\sigma}$, $b = \frac{f_{\sigma'}}{\lambda_{\sigma'}}$ and $c = \frac{f_{\sigma''}}{\lambda_{\sigma''}}$ are Laurent monomials in x,y and z, where f_σ, $f_{\sigma'}$ and $f_{\sigma''}$ belong to σ, σ' and σ'' respectively. Then, an open set $V = \sigma(\Delta_{ij}) \subset G/N$-Hilb$(U)$ is determined by a G/N-graph $\Omega := \{\omega_\tau | \tau \in \mathrm{Irr}(G/N)\} \cong \mathbb{C}[G/N]$ where ω_τ are now monomials in $\mathbb{C}[a,b,c]$. Thus, a point $\mathcal{Z} \in V$ as a G-equivariant module can be written in the form

$$\mathcal{Z} = \{\omega_\tau\Gamma | \tau \in \mathrm{Irr}(G/N)\} = \{\omega_\tau\lambda_\sigma | \tau \in \mathrm{Irr}(G/N), \sigma \in \mathrm{Irr}(N)\} \cong \mathbb{C}[G]$$

In other words, the resulting G-constellations arising from the open set U are obtained by multiplying the building block Γ to the different G/N-graphs Ω.

Example 3.1. Let $G = \frac{1}{6}(1,2,3) = \frac{1}{2}(1,0,1) \times \frac{1}{3}(1,2,0) \cong \mathbb{Z}/6\mathbb{Z}$. Take the normal subgroup in G to be $A = \frac{1}{2}(1,0,1)$. The triangulation of the junior simplex $\Delta = \Delta_1 \cup \Delta_2$ corresponding to A-Hilb$(\mathbb{C}^3) = U_1 \cup U_2$ and the toric coordinates are given in Figure 2.



U_i	$\varepsilon_i, \eta_i, \zeta_i$	A-graph	G/A-action type on U_i
U_1	$\frac{x}{z}, z^2, y$	$\boxed{1\ z}$	$\frac{1}{3}(1,0,2)$
U_2	$\frac{z}{x}, x^2, y$	$\boxed{1\ x}$	$\frac{1}{3}(1,1,1)$

FIGURE 2. Toric fan and coordinates for $\frac{1}{2}(1,0,1)$-Hilb$(\mathbb{C}^3)$.

The action of $G/A \cong \frac{1}{3}(1,2,0)$ on A-Hilb$(\mathbb{C}^3)$ leaves invariant the open sets U_1 and U_2, sending $(\varepsilon_1, \eta_1, \zeta_1) \mapsto (\omega\varepsilon_1, \eta_1, \omega^2\zeta_1)$ and $(\varepsilon_2, \eta_2, \zeta_2) \mapsto (\omega\varepsilon_2, \omega\eta_2, \omega\zeta_2)$ respectively, where ω is a primitive cubic root of unity. Therefore, each of the quotient open sets $U_i/(G/A)$ contains the non-isolated singularities $\frac{1}{3}(1,0,2)$ and $\frac{1}{3}(1,1,1)$ respectively, which we resolve with the crepant resolutions (G/A)-Hilb(U_i). See Figure 3 to see the triangulations of the triangles Δ_1 and Δ_2.

The crepant resolution obtained in this way corresponds to adding to the triangulation $\Delta_1 \cup \Delta_2$ the rest of the points of G-action that we have not used in A-Hilb$(\mathbb{C}^3)$ (in this case $\frac{1}{6}(1,2,3)$, $\frac{1}{6}(2,4,0)$ and $\frac{1}{6}(4,2,0)$ shown in red color) and triangulate every Δ_i according to G/A-Hilb(U_i) for $i = 1,2$. Changing back to the coordinates x, y and z we obtain the triangulation shown in Figure 4.

For $j = 1,2,3$ the basis for the G-constellations of the open set U_{1j} are given by multiplying every basis element in the (G/A)-graphs $\Omega_1 = \{1, \zeta_1, \zeta_1^2\}$, $\Omega_2 = \{1, \zeta_1, \varepsilon_1\}$ and $\Omega_3 = \{1, \varepsilon_1, \varepsilon_1^2\}$ by the building block $\Gamma = \{1, z\}$ coming from the open set U_1. Similarly for the open sets



U_{ij}	Coordinates	G/A-graph
U_{11}	$\frac{\varepsilon_1}{\zeta_1^2}, \zeta_1^3, \eta_1$	$\boxed{1\ \zeta_1\ \zeta_1^2}$
U_{12}	$\frac{\zeta_1^2}{\varepsilon_1}, \frac{\varepsilon_1^2}{\zeta_1}, \eta_1$	$\boxed{1\ \zeta_1\ \varepsilon_1}$
U_{13}	$\varepsilon_1^3, \frac{\zeta_1}{\varepsilon_1^2}, \eta_1$	$\boxed{1\ \varepsilon_1^2\ \varepsilon_1}$



U_{ij}	Coordinates	G/A-graph
U_{21}	$\frac{\varepsilon_2}{\zeta_2}, \frac{\eta_2}{\zeta_2}, \zeta_2^3$	$\boxed{1\ \zeta_2\ \zeta_2^2}$
U_{22}	$\frac{\eta_2}{\varepsilon_2}, \frac{\zeta_2}{\varepsilon_2}, \varepsilon_2^3$	$\boxed{1\ \varepsilon_2\ \varepsilon_2^2}$
U_{23}	$\frac{\varepsilon_2}{\eta_2}, \frac{\zeta_2}{\eta_2}, \eta_2^3$	$\boxed{1\ \eta_2\ \eta_2^2}$

FIGURE 3. $\frac{1}{3}(1,0,2)$-Hilb(U_1) and $\frac{1}{3}(1,1,1)$-Hilb(U_2).



FIGURE 4. Toric fan for $\frac{1}{3}(1,2,0)$-Hilb$(\frac{1}{2}(1,0,1)$-Hilb$(\mathbb{C}^3))$.

$U_{2j} \subset \frac{1}{2}(1,0,1)$-Hilb$(\frac{1}{3}(1,2,0)$-Hilb$(\mathbb{C}^3))$. More precisely, the G-constellations are

$$M_{11} = \{1 \cdot \boxed{1\ z}, \zeta_1 \cdot \boxed{1\ z}, \zeta_1^2 \cdot \boxed{1\ z}\} = \{1, z, y, yz, y^2, y^2z\}$$
$$M_{12} = \{1 \cdot \boxed{1\ z}, \zeta_1 \cdot \boxed{1\ z}, \varepsilon_1 \cdot \boxed{1\ z}\} = \{1, z, y, yz, \frac{x}{z}, x\}$$
$$M_{13} = \{1 \cdot \boxed{1\ z}, \varepsilon_1^2 \cdot \boxed{1\ z}, \varepsilon_1 \cdot \boxed{1\ z}\} = \{1, z, \frac{x^2}{z^2}, \frac{x^2}{z}, \frac{x}{z}, x\}$$
$$M_{21} = \{1 \cdot \boxed{1\ x}, \zeta_2 \cdot \boxed{1\ x}, \zeta_2^2 \cdot \boxed{1\ x}\} = \{1, x, y, xy, y^2, xy^2\}$$
$$M_{22} = \{1 \cdot \boxed{1\ x}, \varepsilon_2 \cdot \boxed{1\ x}, \varepsilon_2^2 \cdot \boxed{1\ x}\} = \{1, x, \frac{z}{x}, z, \frac{z^2}{x^2}, \frac{z^2}{x}\}$$
$$M_{23} = \{1 \cdot \boxed{1\ x}, \eta_2 \cdot \boxed{1\ x}, \eta_2^2 \cdot \boxed{1\ x}\} = \{1, x, x^2, x^3, x^4, x^5\}$$

Let now $A = \frac{1}{3}(1,2,0)$ be the normal subgroup. Then A-Hilb$(\mathbb{C}^3) = V_1 \cup V_2 \cup V_3$ where $V_i \cong \mathbb{C}^3$. The quotient group $G/A \cong \frac{1}{2}(1,0,1)$ produces a $\mathbb{Z}/2\mathbb{Z}$-action on every V_i for $i = 1,2,3$. The resolution of these singularities is translated into the junior simplex $\Delta = \Delta_1 \cup \Delta_2 \cup \Delta_3$ by adding the points $\frac{1}{6}(3,0,3)$ and $\frac{1}{6}(1,2,3)$ and triangulate in the only possible way as in Figure 5. All crepant resolutions of $\mathbb{C}^3/\frac{1}{6}(1,2,3)$ are shown in Figure 6.



	G-constellations
M'_{11}	$\{1, y, y^2, z, yz, y^2z\}$
M'_{12}	$\{1, y, y^2, \frac{x}{y^2}, \frac{x}{y}, x\}$
M'_{21}	$\{1, x, y, \frac{y^2}{x}, y^2, \frac{y^3}{x}\}$
M'_{22}	$\{1, x, y, z, xz, yz\}$
M'_{31}	$\{1, x, x^2, x^3, x^4, x^5\}$
M'_{32}	$\{1, x, x^2, z, xz, x^2z\}$

FIGURE 5. Toric fan $\frac{1}{2}(1,0,1)$-Hilb$(\frac{1}{3}(1,2,0)$-Hilb$(\mathbb{C}^3))$.


$\mathbb{Z}/2\mathbb{Z}$-Hilb($\mathbb{Z}/3\mathbb{Z}$-Hilb($\mathbb{C}^3$))
G-Hilb($\mathbb{C}^3$)
$\mathbb{Z}/3\mathbb{Z}$-Hilb($\mathbb{Z}/2\mathbb{Z}$-Hilb($\mathbb{C}^3$))

Figure 6. Crepant resolutions of $\mathbb{C}^3/G$ with $G = \frac{1}{6}(1,2,3)$.

3.1. **Local coordinates.** The local coordinates of an open set U corresponding to a G-constellation M can be obtained as follows. Let $S := \mathbb{C}[x,y,z]$ and for every $\rho \in \operatorname{Irr}(G)$ consider the Cohen-Macaulay S^G-module $S_\rho := (S \otimes \rho^*)^G$. Taking $M = \bigoplus_{\rho \in \operatorname{Irr}(G)} S_\rho$, the McKay quiver of G is the quiver associated to $\operatorname{End}_{S^G}(M)$.

Let M be a G-constellation in $\mathbb{C}^3$. As a representation of Q, M predetermines the linear independent maps between the vector spaces W_i located at every vertex of $i \in Q$, which give the local coordinates of the corresponding open set. These distinguished linearly independent paths in kQ form *skeleton* of M.

For instance, let $M_{22} = \{1, x, \frac{z}{x}, z, \frac{z^2}{x^2}, \frac{z^2}{x}\}$ be the G-constellation defining the open $U_{22} \in \mathbb{Z}/3\mathbb{Z}$-Hilb$(Y)$ where $Y := \mathbb{Z}/2\mathbb{Z}$-Hilb$(\mathbb{C}^3)$. Let Q be the McKay quiver of G with the usual commutativity relations $xy = yx$, $xz = zx$ and $yz = zy$ (see Figure 7), and consider M_{22} as a representation of Q. Then by choosing the basis element at every vector space $\mathbb{C}_\rho$ to be given by the unique element $\lambda_\rho \in M$, we have that $x \cdot x = a \cdot \frac{z}{x}$, $y \cdot 1 = b \cdot \frac{z}{x}$ and $z \cdot \frac{z^2}{x^2} = c \cdot x$, so that $a = \frac{x^3}{z}$, $b = \frac{xy}{z}$ and $c = \frac{z^3}{x^3}$ which are precisely the local coordinates of $\sigma(\Delta_{22})$. Since after change of basis any nonzero map can be chosen to be 1, the skeleton $sk(U'_{12})$ in this case is formed by the linear maps equal to 1.



Figure 7. McKay quiver for $G = \frac{1}{6}(1,2,3)$ and the open set $U'_{12} \cong \mathbb{C}^3_{a,b,c}$.

Compare with [Ish] Lemma 3.2 for an alternative method when G is an Abelian subgroup in $\mathrm{SL}(3,\mathbb{C})$. We want to emphasize that the method described above and used in this paper is valid for an arbitrary group G.

3.2. **θ-stability in the Abelian case.** Every irreducible representation $\sigma_i \in \operatorname{Irr}(N)$ is fixed so all stabilizers G_i are isomorphic to G/N. The irreducible representations of G are therefore distributed as in Table 3. Since $n_i = d_i^N = d_j^{G/N} = 1$ for all $i = 0, \ldots, p-1$, $j = 0, \ldots, q-1$, we have that $\dim(\rho) = 1$ for $\rho \in \operatorname{Irr}(G)$. The stability condition $\theta \in \Theta$ for which the crepant resolution G/A-Hilb(A-Hilb($\mathbb{C}^3$)) $\cong \mathcal{M}_\theta$ is given also in Table 3.

$$\mathrm{Irr}(G)=\begin{array}{c|c|c|c|c|}
 & \sigma_0 & \sigma_1 & \dots & \sigma_{p-1} \\ \hline
 & \rho_0^0 & \rho_1^0 & \dots & \rho_{p-1}^0 \\ \hline
 & \rho_0^1 & \rho_1^1 & \dots & \rho_{p-1}^1 \\ \hline
 & & \vdots & & \vdots \\ \hline
 & \rho_0^{q-1} & \rho_1^{q-1} & \dots & \rho_{p-1}^{q-1} \\ \hline
\end{array}
\quad
\theta=\begin{array}{|c|c|c|c|}
\sigma_0 & \sigma_1 & \dots & \sigma_{p-1} \\ \hline
-\sum_{i=1}^{p-1}\theta_i^N-\varepsilon\sum_{j=1}^{q-1}\theta_j^{G/N} & \theta_1^N & \dots & \theta_{p-1}^N \\ \hline
-\sum_{i=1}^{p-1}\theta_i^N+\varepsilon\theta_1^{G/N} & \theta_1^N & \dots & \theta_{p-1}^N \\ \hline
\vdots & \vdots & & \vdots \\ \hline
-\sum_{i=1}^{p-1}\theta_i^N+\varepsilon\theta_{q-1}^{G/N} & \theta_1^N & \dots & \theta_{p-1}^N \\ \hline
\end{array}$$

TABLE 3. $\mathrm{Irr}(G)$ and the stability condition θ in terms of θ_N and $\theta_{G/N}$.

Example 3.2. Let us take the group $G=\frac{1}{6}(1,2,3)$ and consider $\frac{1}{3}(1,2,0)$-Hilb$(\frac{1}{2}(1,0,1)$-Hilb$(\mathbb{C}^3))$. The distribution of $\mathrm{Irr}(G)$ and the stability condition θ are shown in Figure 8 where $a,b_1,b_2\in\mathbb{Q}$ are positive numbers and $0<\varepsilon<<1$.

$$\mathrm{Irr}(G)=\begin{array}{|c|c|}\hline
\rho_0^0 & \rho_1^0 \\ \hline
\rho_0^1 & \rho_1^1 \\ \hline
\rho_0^2 & \rho_1^2 \\ \hline
\end{array}
\qquad
\theta:=\begin{array}{|c|c|}\hline
\theta_0 & \theta_1 \\ \hline
\theta_2 & \theta_3 \\ \hline
\theta_4 & \theta_5 \\ \hline
\end{array}
=\begin{array}{|c|c|}\hline
-a-\varepsilon(b_1+b_2) & a \\ \hline
-a+\varepsilon b_1 & a \\ \hline
-a+\varepsilon b_2 & a \\ \hline
\end{array}$$

FIGURE 8. $\mathrm{Irr}(G)$ and stability condition for $\mathbb{Z}/3$-Hilb$(\mathbb{Z}/2$-Hilb$(\mathbb{C}^3))$.

The stability condition is in this case is given clockwise around the McKay quiver. The G-constellations in every open set and the non-zero maps are shown in Figure 9. By checking the subrepresentations in every affine piece we can see that chamber $C\subset\Theta$ is given by

$$\begin{array}{rrr}
\theta_2,\theta_4<0, & \theta_2+\theta_5>0, & \theta_1+\theta_4>0, \\
\theta_3>0, & \theta_4+\theta_5>0, & \theta_0+\theta_1+\theta_3>0.
\end{array}$$

Thus if we take $0<\varepsilon<a/\max\{b_1,b_2\}$ every inequality is satisfied by θ.



FIGURE 9. Constellations for $\mathbb{Z}/3$-Hilb$(\mathbb{Z}/2$-Hilb$(\mathbb{C}^3))$ as representations of Q.

4. NON-ABELIAN EXAMPLES

4.1. **Dihedral groups $D_{2n}\subset\mathrm{SO}(3)$.** The group is generated by

$$D_{2n}:=\left\langle \alpha=\frac{1}{n}(1,-1,0),\beta=\begin{pmatrix}0&1&0\\1&0&0\\0&0&-1\end{pmatrix}\right\rangle$$

and it has order $2n$. The normal subgroup $N := \langle\alpha\rangle$ has n 1-dimensional irreducible representations $\sigma_i(\alpha) = \varepsilon^i$ where ε is a n-th root of unity and $i = 0, \dots, 2n-1$. The action of $G/N = \langle\overline{\beta}\rangle \cong \mathbb{Z}/2\mathbb{Z}$ give the irreducible representations of G as in Table 4 depending on the parity of n.

σ_0	$\{\sigma_1, \sigma_{2m-1}\}$		$\{\sigma_{m-1}, \sigma_{m+1}\}$	σ_m
ρ_0^0	ρ_1^0	$\cdots$	ρ_{m-1}^0	ρ_m^0
ρ_0^1				ρ_n^1

(a)

σ_0	$\{\sigma_1, \sigma_{2m}\}$		$\{\sigma_m, \sigma_{m+1}\}$
ρ_0^0	ρ_1^0	$\cdots$	ρ_m^0
ρ_0^1			

(b)

TABLE 4. $\mathrm{Irr}(D_{2n})$ for (a) $n = 2m$ even, and (b) $n = 2m+1$ odd.

The dimension vectors are ${}^{1}_{1}2\dots2\,{}^{1}_{1}$ and ${}^{1}_{1}2\dots2$ respectively. In this case, as well as for any subgroup $G \subset \mathrm{SO}(3)$, the fibre over the origin $f^{-1}(0)$ of any crepant resolution $f : Y \to \mathbb{C}^3/G$ has dimension 1. See [GNS], [NdCS].

The open cover of N-Hilb$(\mathbb{C}^3)$ is given by n open sets $\bigcup_{j=1}^n U_j$, covering $n-1$ rational curves E_i for $i = 1, \dots, n-1$. The action G/N on N-Hilb$(\mathbb{C}^3)$ identifies U_i and U_{n-i+1} for $i = 1, \dots, n$.

If $n = 2m$ is even then E_m is fixed by G/N having two fixed lines L_+ and L_- crossing transversally the $\mathbb{P}^1$ covered by U_m and U_{m+1}. They give rise to E_+ and E_- respectively. If $n = 2m+1$ is odd then the open set U_{m+1} is fixed by G/N and there is just one fixed line L, producing the new rational curve E.

Diagonalizing the action of G/N we see that in both cases these singularities are of type $\frac{1}{2}(1,1,0)$. By blowing up these singular lines it follows that the dual graph of the fibre over the origin of the singularity is shown in Figure 10.



(a) (b)

FIGURE 10. Dual graph of $f^{-1}(0)$ for G/N-Hilb(N-Hilb$(\mathbb{C}^3)$) for (a) $n = 2m$ even, and (b) $n = 2m+1$ odd. The curve $\widetilde{E}_i$ denote the strict transform of E_i and the numbers denote the degree of the normal bungle at every curve.

In this case we can see the contrast with G-Hilb$(\mathbb{C}^3)$ where in the case n even the graph is different, whereas in the case n odd the difference resides in the degrees of the normal bundles. The fibre over the origin in G-Hilb$(\mathbb{C}^3)$ is shown in Figure 11. See [GNS] and [NdCS] for more details.



(a) (b)

FIGURE 11. Dual graph of $f^{-1}(0)$ for G-Hilb$(\mathbb{C}^3)$ for (a) $n = 2m$ even, and (b) $n = 2m+1$ odd.

We illustrate the construction of G/N-Hilb(N-Hilb$(\mathbb{C}^3)$) in this case by an example. The general case is analogous.

Example 4.1. Let $G = D_{12}$. In this case the McKay quiver with relations (Q, R) is given in Figure 4.1 where the relations R provide the Morita equivalent between $\mathbb{C}Q/\langle R\rangle$ and $S * G$, and are obtained following [BSW].



$$bC = 0, b'C' = 0, cB = 0, c'B' = 0,$$
$$Ca = uB, C'a' = vB', Ac = bu, A'c' = b'v,$$
$$BA = uC, B'B' = vC', ab = cu, a'b' = c'v$$
$$Bb + Cc = dD, B'b' + C'c' = Dd$$
$$Du = vD, ud = dv$$

FIGURE 12. The McKay of D_{12} with relations.

In this case the stability condition given in Definition 2.4 is shown below, where $a_i, b > 0$ for $i = 1, \ldots, 5$ and $0 < \varepsilon << 1$.

$$\theta := \begin{array}{c|c|c|c|}
\sigma_0 & \{\sigma_1, \sigma_5\} & \{\sigma_2, \sigma_4\} & \sigma_3 \\
\hline
-\sum_{i=1}^5 a_i - \varepsilon b & a_1 + a_5 & a_2 + a_4 & a_3 \\
\hline
-\sum_{i=1}^5 a_i + \varepsilon b & & & a_3 \\
\hline
\end{array}$$

We start by considering the gluing of U_1 and U_2 with U_5 and U_6 by the action of G/N. Since U_3 and U_4 contain the fixed part we will treat them separately. We need to consider the N-constellations at the origin of each open chart, so after choosing a basis for every $H^0(gZ)$ for $g \in G/N$ consisting of N-graphs, we obtain the following G-constellations:

N-constellations	G-constellations				Skeleton in $\Bbbk Q$	Open set
$Z_1 = \{1, x, x^2, x^3, x^4, x^5\}$	1	(x, y) (x^5, y^5)	(x^2, y^2) (x^4, y^4)	$x^3 + y^3$		V_1
$gZ_1 = \{1, y, y^2, y^3, y^4, y^5\}$	α			$x^3 - y^3$		
$Z_2 = \{1, y, x, x^2, x^3, x^4\}$	1	(x, y) $(\alpha x, \alpha y)$	(x^2, y^2) (x^4, y^4)	$x^3 + y^3$		V_2
$gZ_2 = \{1, x, y, y^2, y^3, y^4\}$	α			$x^3 - y^3$		

In the skeleton of the open sets the two vertical dots in the two middle vertices of the McKay quiver denote the two linearly independent vectors e_1 and e_2 in the corresponding vector space at that vertex.

With a similar calculation as in Section 3.1 we can calculate the local coordinates to obtain $V_1 \cong \operatorname{Spec} \mathbb{C}[\frac{z(x^3+y^3)}{x^3-y^3}, \frac{xy}{(x^3+y^3)^2}, (x^3+y^3)^2]$, $V_2 \cong \operatorname{Spec} \mathbb{C}[\frac{z(x^3+y^3)}{x^3-y^3}, \frac{x^2y^2}{(x^3+y^3)^2}, \frac{(x^3+y^3)^2}{xy}]$ and $\alpha = \frac{x^3-y^3}{x^3+y^3}$.

Let us now consider the G-constellations arising from the blowup of the fixed lines L_+ and L_-. In the open set $U_3 \cong \mathbb{C}^3_{a,b,c}$ with $a = x^4/y^2$, $b = y^3/x^3$ and $c = z$, every N-cluster is given by the ideal $I_{a,b,c} = (x^4 - ay^2, y^3 - bx^3, xy - ab, z - c)$. Then the lines $L_\pm$ are defined by $b = \pm 1$, $c = 0$, which means that the ideals defining the lines are $I_{L_\pm} = (x^4 - ay^2, y^3 \mp x^3, xy \mp a, z)$. Therefore we can choose as basis for the N-constellations at these lines the N-graphs $\Gamma_\pm = \{1, x, y, x^2, y^2, x^3 \pm y^3\}$, which are invariant under the action of G/N.

For the line L_+ we have $b = 1$. By the change of coordinate $b^+ = 1 - b$ we have that the action of G/N as $\frac{1}{2}(1, 1, 0)$ is defined on $\mathbb{C}^3_{A,B,C}$ where $A = c$, $B = \frac{2-b^+}{b^+}$, $C = a(b^+)^2$. This implies that the rational curve E_+ is covered by the open sets $V_3 \cong \mathbb{C}^3_{B^2,A/B,C}$, $V_6 \cong \mathbb{C}^3_{A^2,B/A,C}$, and E^+ is given by the ratio $(A : B) = (z(x^3 - y^3) : x^3 + y^3)$. In terms of G-constellations, the calculation is the same as in the Abelian case, where V_3 and V_6 are the open cover of $\frac{1}{2}(1,1,0)$-Hilb$(\mathbb{C}^3_{A,B,C})$. See Figure 13. In the case of V_3 for instance, we have the nonzero maps coming from the N-constellations Γ_+ and $\Gamma_+ \cdot B$ (generated from ρ_0^0 and ρ_0^1 respectively) and the extra arrow comes from the fact that $(x^3+y^3)B = (x^3-y^3) + \frac{4x^3y^3}{x^3-y^3}$, which is induced by the G/N-equivariant map $\Gamma_+ \to B\Gamma_+$.

G-constellations

$\Gamma_+ \cdot \{1, B\}$:

1	(x,y)	(x^2,y^2)	x^3+y^3
B	(xB,yB)	(x^2B,y^2B)	$(x^3+y^3)B$

$\Gamma_+ \cdot \{1, z\}$:

1	(x,y)	(x^2,y^2)	x^3+y^3
z	(xz,yz)	(x^2z,y^2z)	$(x^3+y^3)z$

$\Gamma_- \cdot \{1, B'\}$:

1	(x,y)	(x^2,y^2)	$(x^3-y^3)B'$
B'	(xB',yB')	(x^2B',y^2B')	x^3-y^3

$\Gamma_+ \cdot \{1, z\}$:

1	(x,y)	(x^2,y^2)	x^3-y^3
z	(xz,yz)	(x^2z,y^2z)	$z(x^3-y^3)$



Open set

V_3

V_6

V_4

V_5

FIGURE 13. G-constellations at the open sets covering the exceptional curves E^+ and E^-, where $B = \frac{x^3+y^3}{x^3-y^3}$ and $B' = \frac{x^3-y^3}{x^3+y^3}$.

Similarly, for the line L_+ we have $b = -1$. By the change of coordinate $b^+ = 1 - b$, the action of G/N as $\frac{1}{2}(1,1,0)$ is defined on $\mathbb{C}^3_{A',B',C'}$ where $A' = c$, $B' = \frac{b^- - 2}{b^-}$, $C' = a(b^-)^2$. This implies that the rational curve E_- is covered by the open sets $V_4 \cong \mathbb{C}^3_{B'^2,A'/B',C'}$, $V_5 \cong \mathbb{C}^3_{A'^2,B'/A',C'}$, and E^- is given by the ratio $(A' : B') = (z(x^3+y^3) : x^3 - y^3)$. In terms of G-constellations, the calculation is the same as before but taking now Γ_- instead of Γ_+. See Figure 13.


ρ_0^0
ρ_0^1
ρ_1^0
ρ_2^0
ρ_3^0
ρ_3^1
$(1,0)$
$\left(\begin{smallmatrix} 0 & a \\ aC' & 0 \end{smallmatrix}\right)$
$\left(\begin{smallmatrix} 0 & a \\ aC' & 0 \end{smallmatrix}\right)$
$\left(\begin{smallmatrix} 1 \\ C' \end{smallmatrix}\right)$
$\left(\begin{smallmatrix} K \\ 0 \end{smallmatrix}\right)$
$\left(\begin{smallmatrix} 1 & 0 \\ 0 & 1 \end{smallmatrix}\right)$
$(-c', c')$
aC'
z
a
aC'
$(0,1)$
$\left(\begin{smallmatrix} K & 0 \\ 0 & K \end{smallmatrix}\right)$
$\left(\begin{smallmatrix} 1 \\ 1 \end{smallmatrix}\right)$
$\left(\begin{smallmatrix} 0 \\ K \end{smallmatrix}\right)$
$(c'C', -c')$

FIGURE 14. Open set $V_3 \subset \mathcal{M}_\theta$.

The skeletons provide the choices of nonzero variables in the representation space rep(Q). For example, the case V_3 can we choose $c = (1,0)$, $b = (0,1)$, $d = \mathrm{Id}$, $C_1 = 1$, $B = \left(\begin{smallmatrix} 1 \\ 1 \end{smallmatrix}\right)$ and use the relations we obtain the representation space of $V_3 \subset \mathcal{M}_\theta$ shown in Figure 14 where $K = c'(C'-1)$. Thus $V_3 \cong \mathbb{C}^3_{a,c'C'}$ where $a = \frac{z(x^6+y^6)}{x^6-y^6}$, $c' = \frac{(x^6+y^6)^2}{x^2y^2}$ and $C' = \frac{(x^6-y^6)^2}{(x^6+y^6)^2}$.

4.2. **D_8-Hilb($\frac{1}{3}(1,2,0)$-Hilb($\mathbb{C}^3$)).** Let G be the group $N \rtimes G/N \subset \mathrm{SL}(3,\mathbb{C})$ where N is generated by $g = \frac{1}{3}(1,2,0)$ and $G/N \cong D_8 \subset \mathrm{SO}(3)$ is generated by $\alpha = \frac{1}{4}(1,3,0)$ and $\beta = \left(\begin{smallmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{smallmatrix}\right)$. Let

$\operatorname{Irr}(N) = \{\sigma_0, \sigma_1, \sigma_2\}$ where $\sigma_i(g) = \omega^i$ for $i = 0, 1, 2$ and ω a primitive cube root of unity. In this case we have $\operatorname{Orb}(\sigma_0) = \sigma_0$ and $\operatorname{Orb}(\sigma_1) = \{\sigma_1, \sigma_2\}$ with stabilizers $G_0 = G/N \cong D_8$ and $G_1 = \langle\alpha\rangle \cong \mathbb{Z}/4\mathbb{Z}$. Therefore $\operatorname{Irr} G_0 = \{\tau_0, \tau_1, \tau_2, \tau_3, \tau_4\}$ where $\dim(\tau_i) = 1$ for $i = 0, \dots, 3$ and $\dim(\tau_4) = 2$. The irreducible representations of G are shown in Table 5.

σ_0	$\{\sigma_1, \sigma_2\}$
ρ_0^0	ρ_1^0
ρ_0^1	ρ_1^1
ρ_0^2	ρ_1^2
ρ_0^3	ρ_1^3
ρ_0^4	

, $\mathbf{d} =$

σ_0	$\{\sigma_1, \sigma_2\}$
1	2
1	2
1	2
1	2
2	

, $\theta =$

σ_0	$\{\sigma_1, \sigma_2\}$
$-a-b-5\varepsilon c$	$a+b$
$-a-b+\varepsilon c$	$a+b$
$-a-b+\varepsilon c$	$a+b$
$-a-b+\varepsilon c$	$a+b$
$-2a-2b+\varepsilon c$	

TABLE 5. Irreducible representations of G with their dimensions and the stability condition for G/N-Hilb(N-Hilb($\mathbb{C}^3$)) with $a, b, c > 0$ and $0 < \varepsilon << 1$.

The McKay quiver is shown in Figure 15 which coincides with the Mckay quiver of $D_{24} \subset \mathrm{SO}(3)$ as in 4.1 since $G \cong D_{24} = \langle \frac{1}{12}(1, 11, 0), \beta \rangle$.



FIGURE 15. The McKay quiver of G.

The scheme N-Hilb($\mathbb{C}^3$) can be covered by 3 open sets U_i for $i = 1, 2, 3$ with the corresponding distinguished N-constellations $Z_1 = \{1, x, x^2\}$, $Z_2 = \{1, x, y\}$ and $Z_3 = \{1, y, y^2\}$. By the action of $G/N \cong D_8$ the open set U_2 is fixed while U_1 and U_3 are identified.

The fixed open set U_2 has coordinates $a = \frac{x^2}{y}$, $b = \frac{y^2}{x}$ and $c = z$. If we denote by $(+) := a^2 + b^2$ and $(-) := a^2 - b^2$, we have that G/N-Hilb($\mathbb{C}^3_{a,b,c}$) is covered by 5 open sets given by the following G/N constellations:

$$\Gamma_1 = \{1, a, b, (+), (-), a(+), b(+), -(+)(-)\}$$
$$\Gamma_2 = \{1, a, b, (+), (-), a(+), b(+), c\}$$
$$\Gamma_3 = \{1, a, b, (+), (-), ac, -bc, c\}$$
$$\Gamma_4 = \{1, a, b, c(-), (-), ac, -bc, c\}$$
$$\Gamma_5 = \{1, a, b, (+), c(+), ac, -bc, c\}$$

We obtain in this way the open sets V_i given by the G-constellations $\Gamma_i \cdot Z_2$, for $i = 1, \dots, 5$ shown in Figure 16.

In the case of the orbit $\{U_1, U_3\}$ we have $U_1 \cong \mathbb{C}^3_{d,e,f}$ with coordinates $d = x^3$, $e = y/x^2$, $f = z$, and $U_3 \cong \mathbb{C}^3_{d',e',f'}$ with coordinates $d' = y^3$, $e' = x/y^2$, $f' = z$. In each of the open sets there exist a fixed line with stabilizer subgroup $G_1 = \langle\alpha\rangle \cong \mathbb{Z}/4\mathbb{Z}$. This implies that we have to consider the G_1-graphs

$$\Omega_1 = \{1, c, c^2, c^3\}, \Omega_2 = \{1, c, c^2, d\}, \Omega_3 = \{1, c, d, d^2\}, \Omega_4 = \{1, d, d^2, d^3\}$$

in U_1, and

$$\Omega_1' = \{1, c', c'^2, c'^3\}, \Omega_2' = \{1, c', c'^2, d'\}, \Omega_3 = \{1, c', d', d'^2\}, \Omega_4 = \{1, d', d'^2, d'^3\}$$

in U_3. The identification of U_1 and U_3 by β produces the open sets U_i given by the G-constellations $\Omega_i \cdot Z_1 \cup \Omega_i' \cdot Z_3$ for $i = 1, \dots, 4$. See Figure 16.



FIGURE 16. G-constellations for D_8-Hilb($\frac{1}{3}(1,2,0)$-Hilb($\mathbb{C}^3$)).

Let $Y := \frac{1}{3}(1,2,0)$-Hilb$(\mathbb{C}^3)$, Then the exceptional fibre over the origin $\pi^{-1}(0)$ of the crepant resolution $\pi : Y \to \mathbb{C}^3/N$ consist of two $(-2,0)$-rational curves intersecting in one point. The action of G/N interchange these two curves, producing in $Y/(G/N)$ a single rational curve E with singularities of types $\frac{1}{4}(1,3,0)$ and D_8 at 0 and ∞ respectively. The fibre $\phi^{-1}(0)$ of the crepant resolution $\phi : G/N$-Hilb$(Y) \to \mathbb{C}^3/G$ is therefore given by the following dual graph



where E_i are covered by U_i for $i = 1, \dots, 3$, and F_j are covered by V_j for $j = 1, \dots, 4$, and $\widetilde{E}$ is the strict transform of E.

4.3. **Trihedral group of order 12.** Let G be the group generated by $N := \langle \frac{1}{2}(1,1,0), \frac{1}{2}(1,0,1) \rangle$ and $T := \left(\begin{smallmatrix} 0&1&0 \\ 0&0&1 \\ 1&0&0 \end{smallmatrix}\right)$. The Abelian normal subgroup $N \cong \mathbb{Z}/2 \times \mathbb{Z}/2\mathbb{Z}$ with $\mathrm{Irr}(N) = \{\sigma_0, \sigma_1, \sigma_2, \sigma_3\}$ induce the irreducible representations of G as in Table 6. The McKay quiver with relations (Q, R) is given in Figure 17.

By acting first on $\mathbb{C}^3_{x,y,z}$ with N we have that N-Hilb$(\mathbb{C}^3)$ is given by 4 affine open sets U_i, $i = 1, \dots, 4$. The N-constellations at each of these open sets are

$$Z_1 = \{1, x, y, xy\}, Z_2 = \{1, y, z, yz\}, Z_3 = \{1, x, z, xz\}, Z_4 = \{1, x, y, z\}.$$

σ_0	$\{\sigma_1,\sigma_2,\sigma_3\}$
ρ_0^0	ρ_1
ρ_0^1	
ρ_0^2	

, $\mathbf{d}=$

σ_0	$\{\sigma_1,\sigma_2,\sigma_3\}$
1	3
1	
1	

, $\theta=$

σ_0	$\{\sigma_1,\sigma_2,\sigma_3\}$
$-\sum a_i-2\varepsilon b$	$a_1+a_2+a_3$
$-\sum a_i+\varepsilon b$	
$-\sum a_i+\varepsilon b$	

TABLE 6. Irreducible representations of G with their dimensions and the stability condition for G/N-Hilb(N-Hilb($\mathbb{C}^3$)) with $a_i,b>0$ for $i=1,2,3$ and $0<\varepsilon<<1$.



$$uA=vA,\ au=av$$
$$uB=\omega vB,\ bu=\omega bv$$
$$uC=\omega^2vC,\ cu=\omega^2cv$$
$$Aa+\omega Bb+\omega^2Cc=u^2$$
$$Aa+\omega^2Bb+\omega Cc=v^2$$

FIGURE 17. McKay quiver of G with relations.

The action of $G/N=\langle\bar{T}\rangle\cong\mathbb{Z}/3\mathbb{Z}$ identifies the open sets U_1, U_2 and U_3, and fixes U_4, inducing the corresponding action on the N-constellations. The orbit $\{U_1,U_2,U_3\}$ give rise to the open set $V_1\subset G/N$-Hilb(N-Hilb($\mathbb{C}^3$)) shown in 18.

N-constellations

$Z_1=\{1,x,y,xy\}\in U_1$
$T\cdot Z_1=\{1,y,z,yz\}\in U_2$
$T^2\cdot Z_1=\{1,x,z,xz\}\in U_3$

Splitting into Irr(G)

1	(x,y,z)
λ	(xy,xz,yz)
μ	(x,y,z)

Skeleton in $\Bbbk Q$



Open set

V_1

FIGURE 18. G-constellation arising from the orbit $\{U_1,U_2,U_3\}$.

It follows form the same method as Section 3.1 that $\lambda=\frac{R_2}{R_0}$ and $\mu=\frac{R_1}{R_0}$, where $R_0:=y^2z^2+x^2z^2+x^2y^2$, $R_1:=y^2z^2+\omega x^2z^2+\omega^2x^2y^2$ and $R_2:=y^2z^2+\omega^2x^2z^2+\omega x^2y^2$. The local coordinates of this open set are written at the end of the section.

The remaining case is the fixed N-constellation Z_4 in U_4. The open set $U_4\cong\mathbb{C}^3_{a,b,c}$ has coordinates $a=\frac{yz}{x}$, $b=\frac{xz}{y}$ and $c=\frac{xy}{z}$ and we have $T(a)=b$, $T(b)=c$ and $T(c)=a$. On the other hand, diagonalizing the action of $T\cong\mathbb{Z}/3\mathbb{Z}$, we can consider it to be of type $\frac{1}{3}(1,2,0)$ on $\mathbb{C}^3_{\alpha,\beta,\gamma}$ with the new coordinates $\alpha=a+\omega^2b+\omega c$, $\beta=a+\omega b+\omega^2c$ and $\gamma=a+b+c$. That is,

$$\alpha=\frac{f_1}{f_3},\beta=\frac{f_2}{f_3},\gamma=\frac{f_0}{f_3}$$

where $f_0:=x^2+y^2+z^2$, $f_1:=x^2+\omega^2y^2+\omega z^2$, $f_2:=x^2+\omega y^2+\omega^2z^2$, $f_3:=xyz$ and ω is a primitive cube root of unity.

The situation is therefore identical to the Abelian case, so we need to consider the distinguished G/N-constellations in $\frac{1}{3}(1,2,0)$-Hilb($\mathbb{C}_{\alpha,\beta,\gamma}$), namely $\Gamma_1=\{1,\alpha,\alpha^2\}$, $\Gamma_2=\{1,\alpha,\beta\}$, $\Gamma_3=\{1,\beta,\beta^2\}$. They give rise to the G-constellations $Z_4\cdot\Gamma_i$ and the following skeletons:

G-constellations — Skeleton in $\Bbbk Q$ — Open set

$Z_4\cdot\{1,\alpha,\alpha^2\}$:

1	(x,y,z)
α	$(\alpha,\alpha x,\alpha y)$
α^2	$(\alpha^2,\alpha^2 x,\alpha^2 y)$

V_2

$Z_4\cdot\{1,\alpha,\beta\}$:

1	(x,y,z)
α	$(\alpha,\alpha x,\alpha y)$
β	$(\beta,\beta x,\beta y)$

V_3

$Z_4\cdot\{1,\beta^2,\beta\}$:

1	(x,y,z)
β^2	$(\beta^2,\beta^2 x,\beta^2 y)$
β	$(\beta,\beta x,\beta y)$

V_4



FIGURE 19. G-constellations arising from the non-isolated $\frac{1}{3}(1,2,0)$ line.

It can be checked that the matrices giving the open sets $V_i\subset\mathcal{M}_\theta(Q,R)$ are the following:

$$V_1:\quad a=(1,0,0),b=(0,0,1),c=(1,c_2,1),A=\begin{pmatrix}C_1{+}C_3\\ -c_2C_1A_1{+}c_2B_1C_3\\ C_1(c_2^2A_1{-}1)\end{pmatrix},B=\begin{pmatrix}B_1\\0\\A_1\end{pmatrix},C=\begin{pmatrix}C_1\\0\\B_1(c_2^2C_1{-}1)\end{pmatrix},$$
$$u=\begin{pmatrix}0&1&0\\ C_3{-}\omega C_1&\omega^2c_2C_1&\omega^2C_1{+}\omega B_1\\ \omega^2c_2C_3&-\omega^2(c_2^2C_1{-}1)&-\omega^2c_2C_1\end{pmatrix},v=\begin{pmatrix}0&1&0\\ -\omega^2C_1{+}C_3&\omega c_2C_1&\omega C_1{+}\omega^2B_1\\ \omega c_2C_3&-\omega(c_2^2C_1{-}1)&-\omega c_2C_1\end{pmatrix}.$$

$$V_2:\quad a=(1,0,0),b=(B_1-b_3^2B_3,1,b_3),c=(0,0,1),A=\begin{pmatrix}b_1B_1{+}b_3B_3\\ B_3{-}B_1A_1\\ (b_1b_3{+}1)B_1B_3\end{pmatrix},B=\begin{pmatrix}B_1\\0\\B_3\end{pmatrix},C=\begin{pmatrix}1\\0\\A_1\end{pmatrix},$$
$$u=\begin{pmatrix}0&1&0\\ -\omega^2b_1B_1{+}b_3B_3&\omega B_1&\omega^2{+}\omega b_3B_1\\ \omega B_3&-\omega b_3B_3&-\omega B_1\end{pmatrix},v=\begin{pmatrix}0&1&0\\ -\omega b_1B_1{+}b_3B_3&\omega^2B_1&\omega{+}\omega^2b_3B_1\\ \omega^2B_3&-\omega^2b_3B_3&-\omega^2B_1\end{pmatrix}.$$

$$V_3:\quad a=(1,0,0),b=(0,0,1),c=(c_1,1,1),A=\begin{pmatrix}c_1C_1{+}C_3\\ -C_1A_1{+}B_1C_3\\ C_1(A_1{-}c_1^2)\end{pmatrix},B=\begin{pmatrix}B_1\\0\\A_1\end{pmatrix},C=\begin{pmatrix}C_1\\0\\B_1(C_1{-}c_1)\end{pmatrix},$$
$$u=\begin{pmatrix}0&1&0\\ C_3{-}\omega C_1c_1&\omega^2C_1&\omega^2C_1{+}\omega B_1\\ \omega^2(A_1{-}C_1c_1)&-\omega^2(C_1{-}c_1)&-\omega^2C_1\end{pmatrix},v=\begin{pmatrix}0&1&0\\ C_3{-}\omega^2C_1c_1&\omega C_1&\omega C_1{+}\omega^2B_1\\ \omega(A_1{-}C_1c_1)&-\omega(C_1{-}c_1)&-\omega C_1\end{pmatrix}.$$

$$V_4:\quad a=(1,0,0),b=(0,0,1),c=(C_1-c_3^2C_3,1,c_3),A=\begin{pmatrix}c_1C_1{+}C_3\\ C_3{-}C_1A_1\\ (c_1c_3{+}1)C_1C_3\end{pmatrix},B=\begin{pmatrix}1\\0\\A_1\end{pmatrix},C=\begin{pmatrix}C_1\\0\\C_3\end{pmatrix},$$
$$u=\begin{pmatrix}0&1&0\\ -\omega C_1^2{+}\omega c_3^2C_1C_3{+}c_3C_3&\omega^2C_1&\omega{+}\omega^2c_3C_1\\ \omega^2C_3&-\omega^2c_3C_3&-\omega^2C_1\end{pmatrix},v=\begin{pmatrix}0&1&0\\ -\omega^2C_1^2{+}\omega^2c_3^2C_1C_3{+}c_3C_3&\omega C_1&\omega^2{+}\omega c_3C_1\\ \omega C_3&-\omega c_3C_3&-\omega C_1\end{pmatrix}.$$

Using the quiver between the semi-invariants we can compute the local coordinates at every open set as in the previous examples obtaining:

$$\begin{aligned}
V_1&=\mathbb{C}^3_{B_1,c_2,C_1}=\mathbb{C}[-f_1R_0/R_2,-\sqrt{3}f_3/R_0,-f_2R_0/R_1],\\
V_2&=\mathbb{C}^3_{b_3,B_1,B_3}=\mathbb{C}[-R_1/(\sqrt{3}f_2f_3),\sqrt{3}f_1f_3/R_2,\sqrt{3}f_2^2f_3/R_2],\\
V_3&=\mathbb{C}^3_{B_1,c_1,C_1}=\mathbb{C}[-\sqrt{3}f_1f_3/R_2,-R_0/(\sqrt{3}f_3),\sqrt{3}f_2f_3/R_1],\\
V_4&=\mathbb{C}^3_{c_3,C_1,C_3}=\mathbb{C}[R_2/(\sqrt{3}f_1f_3),\sqrt{3}f_2f_3/R_1,\sqrt{3}f_1^2f_3/R_1].
\end{aligned}$$

Moreover, the glueing between the different open sets is given as follows:

$$\begin{aligned} V_4 \ni (c_3, C_1, C_3) &\longleftrightarrow (-c_3^{-1}, C_1 - c_3^2 C_3, C_1) \in V_3 \\ V_1 \ni (B_1, c_2, C_1) &\longleftrightarrow (B_1 c_2, c_2^{-1}, c_2 C_1) \in V_3 \\ V_2 \ni (b_3, B_1, B_3) &\longleftrightarrow (-B_1, B_1 - b_3^2 B_3, -b_3^{-1}) \in V_3 \end{aligned}$$

Hence, the fiber of the origin of the quotient space are three rational curves meeting in a point. The dual graph with the appropriate degrees of the normal bundles is the following:



In this case the chamber $C \subset \Theta$ for which G/N-Hilb(N-Hilb($\mathbb{C}^3$)) $\cong \mathcal{M}_C$ is given by the inequalities $\theta_1 + \theta_3 > 0$, $\theta_2 + \theta_3 > 0$ and $\theta_1 + \theta_2 + \theta_3 < 0$. On the other hand, the fibre over 0 in G-Hilb($\mathbb{C}^3$) is given by the dual graph



and the chamber C' for G-Hilb($\mathbb{C}^3$) $\cong \mathcal{M}_{C'}$ is given by $\theta_i > 0$ for $i \neq 0$. See [NdCS] for details.

5. When Hilb of Hilb is Hilb

In this section we study the relation between G-Hilb and G/N-Hilb(N-Hilb). Since both of them can be constructed as moduli spaces of representations of the McKay quiver, we can ask the question of when they are isomorphic as moduli spaces and if not, when they are isomorphic as algebraic varieties.

Considering them as moduli spaces we have that G-Hilb $\cong \mathcal{M}_C$ and G-Hilb(N-Hilb) $\cong \mathcal{M}_{C'}$ for some chambers $C, C' \subset \Theta$, where C' is the chamber containing the parameter in Definition 2.4. Then the problem is to determine which groups G admit a normal subgroup N such that $C = C'$. We give complete answer to this question in the case of $G \subset \mathrm{SL}(2,\mathbb{C})$ in Corollary 5.2 and $G \subset \mathrm{SL}(3,\mathbb{C})$ in Theorem 5.4.

As algebraic varieties, in dimension 2 there is nothing to prove since both of them are minimal resolutions of $\mathbb{C}^2/G$ thus isomorphic. For $G \subset \mathrm{SL}(3,\mathbb{C})$ we give a complete answer when the group G is Abelian by using the method of Craw and Reid [CR] to obtain the triangulation of the junior simplex Δ which corresponds to G-Hilb($\mathbb{C}^3$). As we saw in Section 3 the triangulation for G/N-Hilb(N-Hilb($\mathbb{C}^3$)) is given by using the Craw-Reid method in two steps, first for N and then for G/N. Comparing both triangulations we are able to describe in Theorem 5.6 all possible configurations for G and N for which there is an isomorphism of varieties.

5.1. **As moduli spaces.** Let $G \subset \mathrm{SL}(3,\mathbb{C})$ and $N \lhd G$ be a normal subgroup. With the same notation as in Section 2 let $Y_1 := N$-Hilb($\mathbb{C}^3$) and $Y_2 := G/N$-Hilb(Y_1) with universal families $\mathcal{Z}_1$ and $\mathcal{Z}_2$ respectively, and denote by $\mathcal{U} := p_{02*}(\mathcal{O}_{\mathcal{Z}_1 \times_{Y_1} \mathcal{Z}_2})$ the flat family over Y_2 of G-constellations by the projection $p_{02} : Y_2 \times Y_1 \times \mathbb{C}^3 \longrightarrow Y_2 \times \mathbb{C}^3$.

As we know, G-Hilb($\mathbb{C}^3$) $\cong \mathcal{M}_C$ and $Y_2 \cong \mathcal{M}_{C'}$ are both crepant resolutions of $\mathbb{C}^3/G$ isomorphic to a moduli space of G-constellations for some chambers $C, C' \subset \Theta$. Then,

$$\begin{aligned} C = C' &\iff \exists \text{ a closed subscheme } \mathcal{Z} \subset Y_2 \times \mathbb{C}^3 \text{ such that } \mathcal{U} \cong \mathcal{O}_{\mathcal{Z}} \\ &\iff \phi : \mathcal{Z}_2 \times_{Y_1} \mathcal{Z}_1 \longrightarrow Y_2 \times \mathbb{C}^3 \text{ is a closed immersion} \\ &\implies \phi \text{ is injective on the } \mathbb{C}\text{-valued points.} \end{aligned}$$

If we denote by $Y_1^{G/N}$ the fixed locus of the action of G/N into Y_1, we obtain the following sufficient condition for Y_2 and G-Hilb$(\mathbb{C}^3)$ being isomorphic as moduli spaces.

Lemma 5.1. *If* $\tau_1^{-1}(0) \nsubseteq Y_1^{G/N}$ *then* $C \neq C'$.

Proof. Let $g \in G/N$ and let $y, g(y) \in \tau_1^{-1}(0) \subset Y_1$ two distinct points. Then there exists a G/N-cluster $W \in Y_2$ such that $y, g(y) \in \operatorname{Supp}(W)$. Then $(W, y), (W, g(y)) \in \mathcal{Z}_2$ and $(y, 0), (g(y), 0) \in \mathcal{Z}_1$, which implies that $(W, y, 0), (W, g(y), 0) \in \mathcal{Z}_2 \times_{Y_1} \mathcal{Z}_1$ are two distinct points. But then $\phi(W, y, 0) = \phi(W, g(y), 0) = (W, 0)$ so ϕ is not injective. □

As an immediate consequence of the above condition we can conclude that for any finite subgroup $G \subset \mathrm{SL}(2, \mathbb{C})$ they are never the same moduli space.

Corollary 5.2. *Let* $G \subset \mathrm{SL}(2, \mathbb{C})$ *be a finite subgroup. Then* $C \neq C'$.

Proof. Since G and G/N are subgroups of $\mathrm{SL}(2, \mathbb{C})$ they are small, so the action of G/N on Y_1 has no quasireflections. Then $\dim Y_1^{G/N} = 0$ while $\tau_1^{-1}(0)$ is the union of rational curves. □

Remark 5.3. The previous statement is not true if we take G to be a finite subgroup of $\mathrm{GL}(2, \mathbb{C})$. For example, if $G \cong \frac{1}{2r}(1, 1)$ and $N \cong \frac{1}{2}(1, 1)$ we have that $\tau_1^{-1}(0) = Y_1^{G/N}$ and in fact the chambers of Y_2 and G-Hilb$(\mathbb{C}^2)$ are the same. Moreover, even if we assume G to be small, the action of G/N on Y_1 can have quasireflections and therefore G/N-Hilb(Y_1) may not be minimal.

From now on let $G \subset \mathrm{SL}(3, \mathbb{C})$ be a finite subgroup. As the following theorem shows, the cases when Y_2 and G-Hilb$(\mathbb{C}^3)$ coincide are very few.

Theorem 5.4. *If* G *be a finite subgroup of* $\mathrm{SL}(3, \mathbb{C})$. *Then,*

$$C = C' \iff G \cong \frac{1}{2r}(1, 1, 2r-2) \text{ and } N \cong \frac{1}{2}(1, 1, 0)$$

Proof. Recall that if $G \subset \mathrm{SL}(3, \mathbb{C})$ then $\dim Y_1^{G/N} \leq 1$. Therefore if $C = C'$ then by Lemma 5.1 we must have $\dim(\tau_1^{-1}(0)) \leq 1$. Moreover, since the $\tau^{-1}(0)$ is connected it must consist of a single curve. Indeed, if we have more than one curve in $\tau^{-1}(0)$ fixed by G/N then at any intersection point of two curves a 2-dimensional subspace of the tangent space is fixed, thus $\dim Y_1^{G/N} > 1$ a contradiction.

Therefore $N \cong \mathbb{Z}/2\mathbb{Z}$ and we can suppose it to be isomorphic to $\frac{1}{2}(1, 1, 0)$. Then N-Hilb$(\mathbb{C}^3) \cong U_1 \cup U_2$ where $U_1 \cong \mathbb{C}^3_{x^2, z, \frac{y}{x}}$, $U_2 \cong \mathbb{C}^3_{y^2, z, \frac{x}{y}}$ and $\tau^{-1}(0) = E \cong \mathbb{P}^1$ with coordinates $x : y$. After extending the action of G/N on $\mathbb{C}[x, y, z]^N$ naturally into $\mathbb{C}(x, y, z)^N$, we have that G/N fixes E if $g(\frac{x}{y}) = \frac{x}{y}$ for all $g \in G/N$. In other words, g as al element of G can be written in the form $\left(\begin{smallmatrix} \varepsilon & 0 & a \\ 0 & \varepsilon & b \\ 0 & 0 & \varepsilon^{n-2} \end{smallmatrix}\right)$ with $a, b \in \mathbb{C}$ and ε a primitive n-th root of unity. But the group N is normal in G, so g must commute with any element in N. This implies that $a = b = 0$ and since G contains $\frac{1}{2}(1, 1, 0)$ as a subgroup n has to be an even number. Thus $G \cong \frac{1}{2r}(1, 1, 2r-2)$ for some $r > 1$.

Conversely, if the group is of the form $G \cong \frac{1}{2r}(1, 1, 2r-2)$ and $N \cong \frac{1}{2}(1, 1, 0)$, then by the construction of G-constellations in the Abelian case of Section 3, we see that the elements ω_τ for $\tau \in \operatorname{Irr}(G/N)$ are not Laurent monomials. More precisely, for U_1 we have $\omega_{\tau_0} = 1$, $\omega_{\tau_1} = x^2$ and for U_2 we have $\omega_{\tau_0} = 1$, $\omega_{\tau_1} = y^2$, so there are no Laurent monomials in the G-constellations $\mathcal{Z}$ of G/N-Hilb$(N$-Hilb$(\mathbb{C}^3))$. This means that they are precisely the G-graphs of G-Hilb$(\mathbb{C}^3)$, thus the chambers are the same. □

5.2. **As varieties.** In this section we assume that $G \subset \mathrm{SL}(3, \mathbb{C})$ is a finite Abelian subgroup. We use the same notation as in Section 3. We start by recalling the properties of the the triangulation of the junior simplex Δ constructed by Craw and Reid in [CR] corresponding to G-Hilb$(\mathbb{C}^3)$ that we need. By abusing the notation, in what follows we identify G-Hilb$(\mathbb{C}^3)$ with its corresponding triangulation of Δ given by [CR].

A *regular triangle* of side r in Δ is a lattice triangle with $r+1$ points on each edge. In G-Hilb$(\mathbb{C}^3)$ every regular triangle of side r is triangulated with the *regular tesselation*, which is done by drawing $r-1$ parallel lines to the sides of the regular triangle, obtaining r^2 regular triangles of side 1. There are only two types of regular triangles appearing in G-Hilb$(\mathbb{C}^3)$, namely the *corner triangle* and the *meeting of champions*, both shown in Figure 20.



Figure 20. Types of regular triangles: (a) corner triangle and (b) meeting of champions.

In particular, the sides of a regular triangle always extends to one of the vertices e_i. From the construction we can deduce the following properties that we use repeatedly in the rest of the section.

Proposition 5.5. *Consider the triangulation of* Δ *corresponding to G-Hilb*$(\mathbb{C}^3)$*. Then,*

(i) Any line contained in Δ *either passes through one of the vertices* e_i *for* $i=1,2,3$*, or it is contained in a regular triangle.*

(ii) The valency of a vertex v *in* Δ *is either 3, 4, 5 or 6.*

Proof. Part (i) follows from the construction of the triangulation of Δ, and part (ii) form [CR] Corollary 1.4. □

Theorem 5.6. *Let* G *be a finite non-simple Abelian subgroup of* $\mathrm{SL}(3,\mathbb{C})$ *and let* N *be a normal subgroup of* G*, with* $N \neq G, \{1\}$*. Then*

$$G/N\text{-Hilb}(N\text{-Hilb}(\mathbb{C}^3)) \cong G\text{-Hilb}(\mathbb{C}^3)$$

as algebraic varieties if and only if we are in one of the following situations

1. $G/N \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ *for some* $m>1$.
2. $G \cong \frac{1}{r}(1,1,r-2)$ *or* $G \cong \frac{1}{r}(1,r-1,0)$*, i.e.* $\mathbb{C}^3/G$ *has a unique crepant resolution.*
3. $G \cong \frac{1}{2r}(1,a,-a-1)$ *with* $(2r,a)=1$*,* $a^2 \equiv 1$ *(mod 4r) and* $N \cong \frac{1}{2}(1,1,0)$.
4. *There is a subgroup* $G' \subset G$ *containing* N *such that* (G',N) *fits into either* (2) *or* (3) *and* $G/G' \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ *for some* $m>1$.

The proof of the theorem is deduced from Lemmas 5.7 to 5.11. The first lemma shows the biggest family of Abelian groups for which we have an an isomorphism of varieties, and constitutes the case (1) in Theorem 5.6. The rest of the cases are in some sense sporadic.

Lemma 5.7. *If* $G/N \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ *then* G/N*-Hilb*$(N$*-Hilb*$(\mathbb{C}^3)) \cong G$*-Hilb*$(\mathbb{C}^3)$ *as varieties.*

Proof. Assume $G/N \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$. Let $L \supset L' \supset \mathbb{Z}^3$ be the toric lattices for $\mathbb{C}^3/G$ and $\mathbb{C}^3/N$ respectively. Then the assumption implies $L/L' \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$. Since L and L' are generated by elements on the junior simplex, we have decompositions $L = L_0 \oplus \mathbb{Z}e_1$ and $L' = L'_0 \oplus \mathbb{Z}e_1$, where $L_0 = L \cap \mathbb{R}^2_\Delta$ and $L'_0 = L' \cap \mathbb{R}^2_\Delta$.

Then we have $L_0/L_0' \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ for the two-dimensional lattices $L_0 \supset L_0'$, which implies $L_0 = (1/m)L_0'$. So the Newton polygon for G at e_1 is $1/m$ times that for G', thus the triangulations are the same. $\square$

The following lemma justifies the Case (4) in Theorem 5.6 and allows us to obtain isomorphism as varieties between Y_2 and G-Hilb$(\mathbb{C}^3)$ by combining the Cases (1), (2) and (3).

Lemma 5.8. *If there exists a surjection $\phi : G/N \twoheadrightarrow \mathbb{Z}/m\mathbb{Z}\times\mathbb{Z}/m\mathbb{Z}$ for some $m > 1$ then we can replace G by the pullback of* Ker(ϕ) *to G.*

Proof. If there exists a sequence of normal subgroups $G \rhd G' \rhd N$ and $G/G' \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ then we can construct G/N-Hilb(N-Hilb$(\mathbb{C}^3)$) in three steps

$$G/G'\text{-Hilb}(G'/N\text{-Hilb}(N\text{-Hilb}(\mathbb{C}^3)))$$

so by Lemma 5.7 we can take G' instead of G. $\square$

Therefore from now on we assume that no such surjection exists.

Lemma 5.9. *If G/N-Hilb(N-Hilb$(\mathbb{C}^3)$) $\cong$ G-Hilb$(\mathbb{C}^3)$ then there is no regular triangle of side ≥ 2 in N-Hilb$(\mathbb{C}^3)$.*

Proof. Let T be a regular triangle in Δ of side ≥ 2. Then T is triangulated by the regular tesselation and there always exists a triangle $\Delta_i \subset \Delta$ which does not contain any of the vertices of T. Now consider the action of G/N on N-Hilb$(\mathbb{C}^3)$ and the corresponding triangulation on Δ_i. Since the sides of T extend to some vertex e_i, by Proposition 5.5 (i) there cannot be any new line going out from any vertex of Δ_i. This implies that the action of G/N into Δ_i has to be of the form $\mathbb{Z}/m\mathbb{Z}\times\mathbb{Z}/m\mathbb{Z}$, so that Δ_i is triangulated again with the regular tesselation. But then $G/N \cong \mathbb{Z}/m \times \mathbb{Z}/m\mathbb{Z}$ which contradicts our assumption. $\square$

Lemma 5.10. *The triangulation of Δ corresponding to N-Hilb$(\mathbb{C}^3)$ contains only regular triangles of side 1 if and only if $N \cong \frac{1}{r}(1,1,r-2)$, $\frac{1}{r}(1,r-1,0)$ or $\frac{1}{7}(1,2,4)$*

Proof. Notice that there are only regular triangles of side 1 if and only if there exists a unique crepant resolution of $\mathbb{C}^3/N$, namely N-Hilb$(\mathbb{C}^3)$. Indeed, if a flop from N-Hilb$(\mathbb{C}^3)$ exists then there are 4 points in Δ not 3 of them align, and any triangulation of them would contradict Proposition 5.5 (i). Conversely, if there exist a unique crepant resolution there is no parallelogram in the triangulation of Δ, in particular there is no regular triangle of side bigger than one.

Finally notice that the Abelian groups for which there exist a unique crepant resolution are $\frac{1}{r}(1,1,r-2)$, $\frac{1}{r}(1,r-1,0)$ or $\frac{1}{7}(1,2,4)$. This follows from the fact that either all points are contained in a line or $N \cong \frac{1}{7}(1,2,4)$, otherwise we would have 4 points in Δ with not 3 of them align, hence a flop. $\square$

From the three possibilities for N in the previous lemma we can exclude the case $N \cong \frac{1}{7}(1,2,4)$. Indeed, let $\Delta = \bigcup_{i=1}^{7}\Delta_i$ the triangulation corresponding to N-Hilb$(\mathbb{C}^3)$. Then, on the regular triangle with vertices $\frac{1}{7}(1,2,4)$, $\frac{1}{7}(2,4,1)$ and $\frac{1}{7}(4,1,2)$ in the middle of Δ, the next triangulation created by the action of G/N has to be a regular triangle again (otherwise it would contradict Proposition 5.5 (i)), so we are again in the case of Lemma 5.7.

Now consider the case $N \cong \frac{1}{r}(1,1,r-2)$. Then every lattice point $P_j \in \Delta$ distinct from the vertices e_i for $i = 1,2,3$, are on a line L passing through e_3. If we consider the triangulation induced by G/N, by Proposition 5.5 (i) there are no new lines going out any of the points P_j unless $r = 2$ or 3. Therefore, either $G/N \cong \frac{1}{s}(1,1,s-2)$ so that every new point is again on the line L and G is of type $\frac{1}{n}(1,1,n-2)$, or $N \cong \frac{1}{3}(1,1,1)$, or $N \cong \frac{1}{2}(1,1,0)$, or we are in the case of Lemma 5.8.

Similarly, if $N \cong \frac{1}{r}(1,r-1,0)$ then either $G/N \cong \frac{1}{s}(1,s-1,0)$ for some $s|r$ or $N \cong \frac{1}{2}(1,1,0)$. In any case, we are either in the case (2) of the Theorem 5.6 or N has order 2 or 3.

Let $N \cong \frac{1}{3}(1,1,1)$ and let P be the point in the center of the triangulation $\Delta = \bigcup_{i=1}^{3} \Delta_i$ of N-Hilb$(\mathbb{C}^3)$ where the 3 lines L_i from the vertices e_i meet $(i = 1,2,3)$. Now consider the second triangulation produced by G/N. Because of the "meeting of champions" only one of the lines L_i can be extended, so that the final valency of P is at most 4.

If the valency is 3 then every Δ_i has a regular triangle around P. Since any two of them share a vector which generates the 2-dimensional lattice $\mathbb{Z}^2_\Delta$ it follows that they must have the same side, and by the symmetry of the lattice we can conclude that $G/N \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ as in Lemma 5.7. If the valency is 4 then at least 2 of the Δ_i's must have a regular triangle around P, which must be of the same side since they share a generator of the lattice. See Figure below.





But then there exists a subgroup $G' \cong \frac{1}{3r}(1,1,3r-2)$ such that $G/G' \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ with $m > 1$. Indeed, let M be the middle point of e_1 and e_2. Then every regular triangle inside the triangle e_1PM has e_1 as a vertex and the other two vertices lie on the segment PM. Moreover, these regular triangles have the same area. Then we can see that there is a subgroup $G' \cong \frac{1}{3r}(1,1,3r-2)$ such that these regular triangles for G are basic triangles for G'. If these regular triangles are divided into m^2 basic triangles, then $G/G' \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$.

In the case $N \cong \frac{1}{2}(1,1,0)$ we have the following lemma which gives the case (3) in Theorem 5.6 and finishes the proof.

Lemma 5.11. *Let* $N \cong \frac{1}{2}(1,1,0)$ *and* $Y_2 := G/N$*-Hilb*$(N$*-Hilb*$(\mathbb{C}^3))$. *Then*

$$Y_2 \cong G\text{-}Hilb(\mathbb{C}^3) \Longleftrightarrow G \cong \frac{1}{2r}(1,a,-a-1) \text{ with } a^2 \equiv 1 \ (mod\ 4r)$$

Proof. Let $\Delta = \Delta_1 \cup \Delta_2$ the triangulation for N-Hilb$(\mathbb{C}^3)$ and $P := \frac{1}{2}(1,1,0) \in \Delta$. Suppose that we have the isomorphism as varieties. After the action of G/N on Δ, by Proposition 5.5 (i) the triangles Δ_i must contain regular triangles around P. The sides of the regular triangles must pass through some vertex e_i so there are two possible configurations (see Figure 21).

Notice that any line passing through the point P must go to one of the vertices e_i for $i = 1,2,3$ (otherwise, the sides of two regular triangles would intersect), and in particular there exists the diagonal line $L := e_3P$. Then the vectors v_1, v_2 are sides of a basic triangle, therefore they form a basis of the 2-dimensional lattice $\mathbb{Z}^2_\Delta$.

This imply that the lattice is symmetric with respect to the diagonal line L and in particular the regular triangles around P have the same side. Then it follows that the continued fraction $\frac{2r}{a}$ at the vertex e_3 has to be symmetric with respect to the middle entry, and the boundary of the Newton polygon must contain a lattice point in L. This imply that the expansion of $\frac{2r}{2r-a}$ is also symmetric thus if $x^iy^j \in (\mathbb{Z}^2_\Delta)^\vee$ then $x^jy^i \in (\mathbb{Z}^2_\Delta)^\vee$ for any i and j. In other words, we have the condition $a^2 \equiv 1 \pmod{2r}$ in order to have such symmetric Newton polygon. In addition, the vectors v_1 and v_2 being a basis of $\mathbb{Z}^2_\Delta$ imply that if $a^2 \equiv 1 \pmod{4r}$, so the groups G we are looking for are precisely $G \cong \frac{1}{2r}(1,a,-a-1)$ with $a^2 \equiv 1 \pmod{2n}$.

Conversely, if the group is of the form $G \cong \frac{1}{2r}(1,a,-a-1)$ with $a^2 \equiv 1 \pmod{4r}$, then $\{v_1, v_2\}$ and $\{v_1, -v_2\}$ form a basis of the lattice of Δ. It follows that the line L has to be part of the triangulation and again the distribution of points along Δ is symmetric with respect to L, with regular triangles in Δ_1 and Δ_2 around P. The continued fractions at the vertices e_1 and e_2 are



FIGURE 21. The two possible regular tesselations around $P = \frac{1}{2}(1,1,0)$ and the generators v_1, v_2 of the lattice $\mathbb{Z}^2_\Delta$.

the same, and the one for the vertex e_3 is symmetric with respect to the middle term, so by the Craw-Reid method the triangulation of G-Hilb($\mathbb{C}^3$) and G/N-Hilb(N-Hilb($\mathbb{C}^3$)) are the same, and the result follows. □

Example 5.12. In this example we show a group G with two subgroups, with one of them there is an isomorphism as moduli spaces (hence as varieties) between G-Hilb($\mathbb{C}^3$) and G/N-Hilb(Y_1), and with the other an isomorphism as varieties but in different chamber.

Let $G = \frac{1}{6}(1,1,4)$. By taking $N = \frac{1}{2}(1,1,0)$ and let $Y_1 := N$-Hilb($\mathbb{C}^3$). Then we have an isomorphism G-Hilb($\mathbb{C}^3$) $\cong$ G/N-Hilb(Y_1) as moduli spaces of representations of the McKay quiver.

The crepant resolution Y_1 is covered by $U_1 \cong \mathbb{C}^3_{y^2,z,x/y}$ and $U_1 \cong \mathbb{C}^3_{x^2,z,y/x}$, and $\tau_1^{-1}(0)$ consists of the single $\mathbb{P}^1$ joining the two toric fixed points with coordinates $(x:y)$. In every open set the action of G/N is isomorphic to $\frac{1}{3}(1,2,0)$, and we have that $\tau_1^{-1}(0) = Y_1^{G/N}$. The corresponding G-constellations shown in the Figure below. The chamber C for G-Hilb($\mathbb{C}^3$) is given by the inequalities

$$\theta_3 > 0, \theta_5 > 0, \theta_2 + \theta_3 > 0, \theta_4 + \theta_5 > 0, \theta_1 + \theta_3 + \theta_5 > 0, \theta_0 < 0, \theta_0 + \theta_4 < 0$$



U_{11}	$\{1,y,y^2,y^3,y^4,y^5\}$
U_{12}	$\{1,y,y^2,y^3,z,yz\}$
U_{13}	$\{1,y,z,yz,z^2,y^2\}$
U_{21}	$\{1,x,x^2,x^3,x^4,x^5\}$
U_{22}	$\{1,x,x^2,x^3,z,xz\}$
U_{23}	$\{1,y,z,xz,z^2,xz^2\}$

$$\theta := \begin{array}{|c|c|}\hline \theta_0 & \theta_1 \\ \hline \theta_2 & \theta_3 \\ \hline \theta_4 & \theta_5 \\ \hline \end{array} = \begin{array}{|c|c|}\hline -a-\varepsilon(b_1+b_2) & a \\ \hline -a+\varepsilon b_1 & a \\ \hline -a+\varepsilon b_2 & a \\ \hline \end{array}$$

FIGURE 22. Stability condition for $\mathbb{Z}/3$-Hilb($\mathbb{Z}/2$-Hilb($\mathbb{C}^3$)).

For G/N-Hilb(N-Hilb($\mathbb{C}^3$)), according to Definition 2.4 the stability is given in Figure 22 where $a, b \in \mathbb{Q}$ and $0 < \varepsilon << 1$, which is contained in C.

Now take $N = \frac{1}{3}(1,1,1)$. Then we have that G-Hilb($\mathbb{C}^3$) $\cong$ G/N-Hilb(N-Hilb($\mathbb{C}^3$)) are isomorphic as varieties but in different chambers. In this case N-Hilb($\mathbb{C}^3$) has 3 open sets

$$V_1 \cong \mathbb{C}^3_{y^3,\frac{z}{y},\frac{x}{y}}, V_2 \cong \mathbb{C}^3_{x^3,\frac{z}{x},\frac{y}{x}}, V_3 \cong \mathbb{C}^3_{\frac{y}{z},\frac{x}{z},z^3},$$

and $G/N \cong \frac{1}{2}(1,1,0)$ in every open set. Then we obtain the distinguished G-constellations shown below. In this case the stability condition is given in Figure 23, where in particular we have that $\theta_3 < 0$ so the chamber is different than C.

$$\theta := \begin{array}{|c|c|c|}\hline \theta_0 & \theta_1 & \theta_2 \\ \hline \theta_3 & \theta_4 & \theta_5 \\ \hline \end{array} = \begin{array}{|c|c|c|}\hline -a_1 - a_2 - \varepsilon b & a_1 & a_2 \\ \hline -a_1 - a_2 + \varepsilon b & a_1 & a_2 \\ \hline \end{array}$$

FIGURE 23. Stability condition for $\mathbb{Z}/2$-Hilb($\mathbb{Z}/3$-Hilb($\mathbb{C}^3$)).



V_{11}	$\{1, y, y^2, y^3, y^4, y^5\}$
V_{12}	$\{1, y, y^2, \frac{z}{y}, z, yz\}$
V_{21}	$\{1, x, x^2, x^3, x^4, x^5\}$
V_{22}	$\{1, x, x^2, \frac{z}{x}, z, xz\}$
V_{31}	$\{1, z, z^2, \frac{x}{z}, x, xz\}$
V_{32}	$\{1, z, z^2, \frac{y}{z}, y, yz\}$

REFERENCES

[BKR] Tom Bridgeland, Alastair King, and Miles Reid. The McKay correspondence as an equivalence of derived categories. *J. Amer. Math. Soc.*, 14(3):535–554 (electronic), 2001.

[Bri] Tom Bridgeland. Flops and derived categories. *Invent. Math.*, 147(3):613–632, 2002.

[BSW] Raf Bocklandt, Travis Schedler, and Michael Wemyss. Superpotentials and higher order derivations. *J. Pure Appl. Algebra*, 214(9):1501–1522, 2010.

[CI] Alastair Craw and Akira Ishii. Flops of G-Hilb and equivalences of derived categories by variation of GIT quotient. *Duke Math. J.*, 124(2):259–307, 2004.

[CR] Alastair Craw and Miles Reid. How to calculate A-Hilb $\mathbb{C}^3$. In *Geometry of toric varieties*, volume 6 of *Sémin. Congr.*, pages 129–154. Soc. Math. France, Paris, 2002.

[Cra01] Alastair Craw. *The McKay correspondence and representations of the McKay quiver*. PhD thesis, Warwcik, 2001.

[Cra05] Alastair Craw. An explicit construction of the mckay correspondence for a-hilb$\mathbb{C}^3$. *J. Algebra*, 285(2):682–705, 2005.

[Dem] Laurent Demonet. Skew group algebras of path algebras and preprojective algebras. *J. Algebra*, 323(4):1052–1059, 2010.

[GNS] Yasushi Gomi, Iku Nakamura, and Ken-ichi Shinoda. Coinvariant algebras of finite subgroups of SL(3, $\mathbf{C}$). *Canad. J. Math.*, 56(3):495–528, 2004.

[Ish] Akira Ishii. Representation moduli of the McKay quiver for finite Abelian subgroups of SL(3, $\mathbb{C}$). In *Strings and geometry*, volume 3 of *Clay Math. Proc.*, pages 227–237. Amer. Math. Soc., Providence, RI, 2004.

[IU] Akira Ishii and Kazushi Ueda The special McKay correspondence and exceptional collection. *arXiv:1104.2381*, April 2011.

[Ito94] Yukari Ito. Crepant resolution of trihedral singularities. *Proc. Japan Acad. Ser. A Math. Sci.*, 70(5):131–136, 1994.

[Ito95] Yukari Ito. Crepant resolution of trihedral singularities and the orbifold Euler characteristic. *Internat. J. Math.*, 6(1):33–43, 1995.

[IN] Yukari Ito and Hiraku Nakajima. McKay correspondence and Hilbert schemes in dimension three. *Topology*, 39(6):1155–1191, 2000.

[Kin] A. D. King. Moduli of representations of finite-dimensional algebras. *Quart. J. Math. Oxford Ser. (2)*, 45(180):515–530, 1994.

[Nak] Iku Nakamura. Hilbert schemes of Abelian group orbits. *J. Algebraic Geom.*, 10(4):757–779, 2001.

[NdC] Alvaro Nolla de Celis. Dihedral G-Hilb via representations of the McKay quiver. *ArXiv e-prints*, May 2009.

[NdCS] Alvaro Nolla de Celis and Yuhi Sekiya. Flops and mutations for crepant resolutions of polyhedral singularities. *In preparation*, 2011.

[Tha] Michael Thaddeus. Geometric invariant theory and flips. *J. Amer. Math. Soc.*, 9(3):691–723, 1996.

[YY] Stephen S.-T. Yau and Yung Yu. Gorenstein quotient singularities in dimension three. *Mem. Amer. Math. Soc.*, 105(505):viii+88, 1993.

Department of Mathematics, Graduate School of Science, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, 739-8526, Japan
E-mail address: `akira@math.sci.hiroshima-u.ac.jp`

Graduate School of Mathematics, Nagoya University, Chikusa-ku Nagoya 464-8602 Japan
E-mail address: `y-ito@math.nagoya-u.ac.jp`

Graduate School of Mathematics, Nagoya University, Chikusa-ku Nagoya 464-8602 Japan
E-mail address: `anolla@gmail.com`